

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
WO 01/92340 A2

- (51) International Patent Classification: C07K 16/28, A61K 39/395, A61P 19/02, 31/04, A61K 38/17
- (74) Agent: ANDERSON, Kathryn, A.: 51 University Street, Seattle, WA 98101-2936 (US).
- (21) International Application Number: PCT/US01/17094
- (22) International Filing Date: 25 May 2001 (25.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/579,808 26 May 2000 (26.05.2000) US  
09/665,343 19 September 2000 (19.09.2000) US  
09/785,934 15 February 2001 (15.02.2001) US  
09/847,816 1 May 2001 (01.05.2001) US
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NO, NZ, PL, PT, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): IM-MUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101-2936 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): PLUENNEKE, John, D. [US/US]; 5816 Hickory Place, Parkville, MO 64152 (US).

## Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF INTERLEUKIN-4 ANTAGONISTS AND COMPOSITIONS THEREOF

0001 0002 0003 0004 0005 0006 0007 0008 0009 0010 0011 0012 0013 0014 0015 0016 0017 0018 0019 0020 0021 0022 0023 0024 0025 0026 0027 0028 0029 0030 0031 0032 0033 0034 0035 0036 0037 0038 0039 0040 0041 0042 0043 0044 0045 0046 0047 0048 0049 0050 0051 0052 0053 0054 0055 0056 0057 0058 0059 0060 0061 0062 0063 0064 0065 0066 0067 0068 0069 0070 0071 0072 0073 0074 0075 0076 0077 0078 0079 0080 0081 0082 0083 0084 0085 0086 0087 0088 0089 0090 0091 0092 0093 0094 0095 0096 0097 0098 0099 0100 0101 0102 0103 0104 0105 0106 0107 0108 0109 0110 0111 0112 0113 0114 0115 0116 0117 0118 0119 0120 0121 0122 0123 0124 0125 0126 0127 0128 0129 0130 0131 0132 0133 0134 0135 0136 0137 0138 0139 0140 0141 0142 0143 0144 0145 0146 0147 0148 0149 0150 0151 0152 0153 0154 0155 0156 0157 0158 0159 0160 0161 0162 0163 0164 0165 0166 0167 0168 0169 0170 0171 0172 0173 0174 0175 0176 0177 0178 0179 0180 0181 0182 0183 0184 0185 0186 0187 0188 0189 0190 0191 0192 0193 0194 0195 0196 0197 0198 0199 0200 0201 0202 0203 0204 0205 0206 0207 0208 0209 0210 0211 0212 0213 0214 0215 0216 0217 0218 0219 0220 0221 0222 0223 0224 0225 0226 0227 0228 0229 0230 0231 0232 0233 0234 0235 0236 0237 0238 0239 0240 0241 0242 0243 0244 0245 0246 0247 0248 0249 0250 0251 0252 0253 0254 0255 0256 0257 0258 0259 0260 0261 0262 0263 0264 0265 0266 0267 0268 0269 0270 0271 0272 0273 0274 0275 0276 0277 0278 0279 0280 0281 0282 0283 0284 0285 0286 0287 0288 0289 0290 0291 0292 0293 0294 0295 0296 0297 0298 0299 0300 0301 0302 0303 0304 0305 0306 0307 0308 0309 0310 0311 0312 0313 0314 0315 0316 0317 0318 0319 0320 0321 0322 0323 0324 0325 0326 0327 0328 0329 0330 0331 0332 0333 0334 0335 0336 0337 0338 0339 0340 0341 0342 0343 0344 0345 0346 0347 0348 0349 0350 0351 0352 0353 0354 0355 0356 0357 0358 0359 0360 0361 0362 0363 0364 0365 0366 0367 0368 0369 0370 0371 0372 0373 0374 0375 0376 0377 0378 0379 0380 0381 0382 0383 0384 0385 0386 0387 0388 0389 0390 0391 0392 0393 0394 0395 0396 0397 0398 0399 0400 0401 0402 0403 0404 0405 0406 0407 0408 0409 0410 0411 0412 0413 0414 0415 0416 0417 0418 0419 0420 0421 0422 0423 0424 0425 0426 0427 0428 0429 0430 0431 0432 0433 0434 0435 0436 0437 0438 0439 0440 0441 0442 0443 0444 0445 0446 0447 0448 0449 0450 0451 0452 0453 0454 0455 0456 0457 0458 0459 0460 0461 0462 0463 0464 0465 0466 0467 0468 0469 0470 0471 0472 0473 0474 0475 0476 0477 0478 0479 0480 0481 0482 0483 0484 0485 0486 0487 0488 0489 0490 0491 0492 0493 0494 0495 0496 0497 0498 0499 0500 0501 0502 0503 0504 0505 0506 0507 0508 0509 0510 0511 0512 0513 0514 0515 0516 0517 0518 0519 0520 0521 0522 0523 0524 0525 0526 0527 0528 0529 0530 0531 0532 0533 0534 0535 0536 0537 0538 0539 0540 0541 0542 0543 0544 0545 0546 0547 0548 0549 0550 0551 0552 0553 0554 0555 0556 0557 0558 0559 0560 0561 0562 0563 0564 0565 0566 0567 0568 0569 0570 0571 0572 0573 0574 0575 0576 0577 0578 0579 0580 0581 0582 0583 0584 0585 0586 0587 0588 0589 0590 0591 0592 0593 0594 0595 0596 0597 0598 0599 0600 0601 0602 0603 0604 0605 0606 0607 0608 0609 0610 0611 0612 0613 0614 0615 0616 0617 0618 0619 0620 0621 0622 0623 0624 0625 0626 0627 0628 0629 0630 0631 0632 0633 0634 0635 0636 0637 0638 0639 0640 0641 0642 0643 0644 0645 0646 0647 0648 0649 0650 0651 0652 0653 0654 0655 0656 0657 0658 0659 0660 0661 0662 0663 0664 0665 0666 0667 0668 0669 0670 0671 0672 0673 0674 0675 0676 0677 0678 0679 0680 0681 0682 0683 0684 0685 0686 0687 0688 0689 0690 0691 0692 0693 0694 0695 0696 0697 0698 0699 0700 0701 0702 0703 0704 0705 0706 0707 0708 0709 0710 0711 0712 0713 0714 0715 0716 0717 0718 0719 0720 0721 0722 0723 0724 0725 0726 0727 0728 0729 0730 0731 0732 0733 0734 0735 0736 0737 0738 0739 0740 0741 0742 0743 0744 0745 0746 0747 0748 0749 0750 0751 0752 0753 0754 0755 0756 0757 0758 0759 0760 0761 0762 0763 0764 0765 0766 0767 0768 0769 0770 0771 0772 0773 0774 0775 0776 0777 0778 0779 0780 0781 0782 0783 0784 0785 0786 0787 0788 0789 0790 0791 0792 0793 0794 0795 0796 0797 0798 0799 0800 0801 0802 0803 0804 0805 0806 0807 0808 0809 0810 0811 0812 0813 0814 0815 0816 0817 0818 0819 0820 0821 0822 0823 0824 0825 0826 0827 0828 0829 0830 0831 0832 0833 0834 0835 0836 0837 0838 0839 0840 0841 0842 0843 0844 0845 0846 0847 0848 0849 0850 0851 0852 0853 0854 0855 0856 0857 0858 0859 0860 0861 0862 0863 0864 0865 0866 0867 0868 0869 0870 0871 0872 0873 0874 0875 0876 0877 0878 0879 0880 0881 0882 0883 0884 0885 0886 0887 0888 0889 0890 0891 0892 0893 0894 0895 0896 0897 0898 0899 0900 0901 0902 0903 0904 0905 0906 0907 0908 0909 0910 0911 0912 0913 0914 0915 0916 0917 0918 0919 0920 0921 0922 0923 0924 0925 0926 0927 0928 0929 0930 0931 0932 0933 0934 0935 0936 0937 0938 0939 0940 0941 0942 0943 0944 0945 0946 0947 0948 0949 0950 0951 0952 0953 0954 0955 0956 0957 0958 0959 0960 0961 0962 0963 0964 0965 0966 0967 0968 0969 0970 0971 0972 0973 0974 0975 0976 0977 0978 0979 0980 0981 0982 0983 0984 0985 0986 0987 0988 0989 0990 0991 0992 0993 0994 0995 0996 0997 0998 0999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 219

## TITLE

## USE OF INTERLEUKIN-4 ANTAGONISTS AND COMPOSITIONS THEREOF

## BACKGROUND OF THE INVENTION

5 Interleukin-4 (IL-4), previously known as B cell stimulating factor, or BSF-1, was originally characterized by its ability to stimulate the proliferation of B cells in response to low concentrations of antibodies directed to surface immunoglobulin. IL-4 has been shown to possess a far broader spectrum of biological activities, including growth co-stimulation of T cells, mast cells, granulocytes, megakaryocytes, and erythrocytes. In addition, IL-4 stimulates the proliferation of several IL-2-  
10 and IL-3-dependent cell lines, induces the expression of class II major histocompatibility complex molecules on resting B cells, and enhances the secretion of IgE and IgG1 isotypes by stimulated B cells. IL-4 is associated with a TH2-type immune response, being one of the cytokines secreted by TH2 cells.

Murine and human IL-4 have been identified and characterized, including cloning of IL-4  
15 cDNAs and determination of the nucleotide and encoded amino acid sequences. (See Yokota et al., *Proc. Natl. Acad. Sci. USA* 83:5894, 1986; Noma et al., *Nature* 319:640, 1986; Grabstein et al., *J. Exp. Med.* 163:1405, 1986; and U.S. Patent 5,017,691.)

IL-4 binds to particular cell surface receptors, which results in transduction of a biological signal to cells such as various immune effector cells. IL-4 receptors are described, and DNA and  
20 amino acid sequence information presented, in Mosley et al., *Cell* 59:335-348, October 20, 1989 (murine IL-4R); Idzerda et al., *J. Exp. Med.* 171:861-873, March 1990 (human IL-4R); and U.S. Patent 5,599,905. The IL-4 receptor described in these publications is sometimes referred to as IL-4R $\alpha$ .

Other proteins have been reported to be associated with IL-4R $\alpha$  on some cell types, and  
25 to be components of multi-subunit IL-4 receptor complexes. One such subunit is IL-2R $\gamma$ , also known as IL-2R $\gamma_c$ . (See the discussion of IL-4R complexes in Sato et al., *Current Opinion in Cell Biology*, 6:174-179, 1994.) IL-4R $\alpha$  has been reported to be a component of certain multi-subunit IL-13 receptor complexes (Zurawski et al., *J. Biol. Chem.* 270 (23), 13869, 1995; de Vries, *J. Allergy Clin. Immunol.* 102(2):165, August 1998; and Callard et al. *Immunology Today*,  
30 17(3):108, March 1996).

IL-4 has been implicated in a number of disorders, examples of which are allergy and asthma. Studies of biological properties of IL-4 continue, in an effort to identify additional activities associated with this pleiotrophic cytokine, and to elucidate the role IL-4 may play in various biological processes and diseases.

35

## SUMMARY OF THE INVENTION

The present invention provides methods for treating certain conditions induced by IL-4, comprising administering an IL-4 antagonist to a patient afflicted with such a condition. Also provided are compositions for use in such methods, comprising an effective amount of an IL-4

antagonist and a suitable diluent, excipient, or carrier. Endogenous IL-4 may be contacted with an IL-4 antagonist in alternative methods, such as those involving *ex vivo* procedures.

Among the conditions to be treated in accordance with the present invention are septic arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, ulcerative colitis, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, lung disorders in which IL-4 plays a role, conditions in which IL-4-induced epithelial barrier disruption plays a role, disorders of the digestive system in which IL-4 plays a role, allergic reactions to medication, Kawasaki disease, sickle cell crisis, Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, and nephrosis. IL-4 antagonists also find use as adjuvants to allergy immunotherapy and as vaccine adjuvants.

IL-4 antagonists include, but are not limited to, IL-4 receptors (IL-4R), antibodies that bind IL-4, antibodies that bind IL-4R, IL-4 muteins that bind to IL-4R but do not induce a biological response, molecules that inhibit IL-4-induced signal transduction, and other compounds that inhibit a biological effect that results from the binding of IL-4 to a cell surface IL-4R.

Examples of IL-4 receptors are soluble forms of the human IL-4 receptor of SEQ ID NO:2. Particular antibodies provided herein include human monoclonal antibodies generated by procedures involving immunization of transgenic mice. Such human antibodies may be directed against human IL-4 receptor, for example.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGURES 1A-C present the nucleotide sequence of the coding region of a human IL-4 receptor cDNA. The amino acid sequence encoded by the cDNA is presented as well. The cDNA clone was isolated from a cDNA library derived from a human T cell line T22. The encoded protein comprises (from N- to C-terminus) an N-terminal signal peptide, followed by an extracellular domain, a transmembrane region (underlined), and a cytoplasmic domain, as discussed further below. The DNA and amino acid sequences of Figures 1A to 1C are also presented in SEQ ID NOS:1 and 2, respectively.

Figures 2A to 2C depict targeted insertion of a neo cassette into the Sma I site of the  $\mu 1$  exon. The construct was employed in generating transgenic mice, as described in Example 2. Figure 2A is a schematic diagram of the genomic structure of the  $\mu$  locus. The filled boxes represent the  $\mu$  exons. Figure 2B is a schematic diagram of the CrmD targeting vector. The dotted lines denote those genomic  $\mu$  sequences included in the construct. Plasmid sequences are not shown. Figure 2C is a schematic diagram of the targeted  $\mu$  locus in which the neo cassette has been inserted into  $\mu 1$ .

Figures 3A and 3B present the nucleotide sequence of a vector designated pGP1k, as described in Example 3 below. This nucleotide sequence also is presented in SEQ ID NO:4.

Figure 4 presents an alignment of the variable region of the light chain ( $V_L$  region, designated  $V_k$  for variable kappa chain) amino acid sequences for several antibodies raised

against human IL-4R. These antibodies and sequences are described further in examples 6, 8, and 9 below. The first sequence in the alignment is for an antibody designated 1B7. In the sequences for the other antibodies, residues that differ from the residue found at that position in 1B7 are shown, and residues that are identical to the residue found at that position in 1B7 are indicated by "-".

Figure 5 presents an alignment of the variable region of the heavy chain ( $V_H$ ) amino acid sequences for several antibodies raised against human IL-4R. These antibodies and sequences are described further in examples 6, 8, and 9 below. The first sequence in the alignment is for an antibody designated 1B7. In the sequences for the other antibodies, residues that differ from the residue found at that position in 1B7 are shown, and residues that are identical to the residue found at that position in 1B7 are indicated by "-".

### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides methods for treating certain conditions induced by IL-4, and for inhibiting biological activities of interleukin-4 (IL-4) *in vivo*. One method comprises administering an IL-4 antagonist to a patient afflicted with such a condition. Compositions for use in such methods for treating IL-4-induced conditions also are provided.

Among the conditions to be treated in accordance with the present invention are septic/reactive arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, lung disorders in which IL-4 plays a role, conditions in which IL-4-induced epithelial barrier disruption plays a role, disorders of the digestive system in which IL-4 plays a role, including inflammatory bowel disease and other inflammatory conditions in the gastrointestinal tract, allergic reactions to medication, Kawasaki disease, sickle cell disease (including sickle cell crisis), Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, and nephrosis, as described in more detail below. IL-4 antagonists also find use as adjuvants to allergy immunotherapy and as vaccine adjuvants.

IL-4 antagonists that may be employed include those compounds that inhibit a biological activity of IL-4. Biological activity(ies) of IL-4 that are inhibited by an antagonist in accordance with methods provided herein are activities that play a role in the particular disease to be treated.

Suitable antagonists include, but are not limited to, IL-4 receptors (IL-4R), antibodies that bind IL-4, antibodies that bind IL-4R, IL-4 muteins that bind to IL-4R but do not induce biological responses, molecules that inhibit IL-4-induced signal transduction, and other compounds that inhibit a biological effect that results from the binding of IL-4 to a cell surface IL-4R. Examples of such IL-4 antagonists are described in more detail below. Particular embodiments of the invention are directed to novel antibodies, polypeptides, and nucleic acid molecules, as described below. Antibodies provided herein include, but are not limited to, human monoclonal antibodies that bind to human IL-4 receptor, and that function as antagonists of both IL-4 and IL-13.

### Indications

The present invention provides methods comprising administering an IL-4 antagonist to a patient afflicted with any of a number of conditions induced by IL-4. IL-4-induced conditions include conditions caused or exacerbated, directly or indirectly, by IL-4. Other factors or cytokines also may play a role in such conditions, but IL-4 induces or mediates the condition to some degree, i.e., at least in part.

The biological activities of IL-4 are mediated through binding to specific cell surface receptors, referred to as interleukin-4 receptors (IL-4R). IL-4-induced conditions include those arising from biological responses that result from the binding of IL-4 to a native IL-4 receptor on a cell, or which may be inhibited or suppressed by preventing IL-4 from binding to an IL-4 receptor. Conditions that may be treated include, but are not limited to, medical disorders characterized by abnormal or excess expression of IL-4, or by an abnormal host response to IL-4 production. Further examples are conditions in which IL-4-induced antibody production, or proliferation or influx of a particular cell type, plays a role. IL-4-induced disorders include those in which IL-4 induces upregulation of IL-4 receptors or enhanced production of another protein that plays a role in a disease (e.g., another cytokine).

A method for treating a mammal, including a human patient, who has such a medical disorder comprises administering an IL-4 antagonist to the mammal or otherwise contacting endogenous IL-4 with an antagonist, e.g., in an *ex vivo* procedure. Conditions that may be treated in accordance with the present invention include, but are not limited to, septic/reactive arthritis, dermatitis herpetiformis, urticaria (especially chronic idiopathic urticaria), ulcers, gastric inflammation, mucosal inflammation, ulcerative colitis, Crohn's Disease, inflammatory bowel disease, other disorders of the digestive system in which IL-4 plays a role (e.g., IL-4-induced inflammation of part of the gastrointestinal tract), conditions in which IL-4-induced barrier disruption plays a role (e.g., conditions characterized by decreased epithelial barrier function in the lung or gastrointestinal tract), scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, IL-4-induced pulmonary conditions (including those listed below), allergic reactions to medication, Kawasaki disease, sickle cell disease or crisis, Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, nephrosis, pemphigus vulgaris or bullous pemphigoid (autoimmune blistering diseases), and myasthenia gravis (an autoimmune muscular disease). IL-4 antagonists also find use as adjuvants to allergy immunotherapy and as vaccine adjuvants, especially when directing the immune response toward a TH1 response would be beneficial in treating or preventing the disease in question.

### Septic/reactive arthritis

IL-4 antagonists may be employed in treating septic arthritis, which also is known as reactive arthritis or bacterial arthritis. Septic arthritis can be triggered by (result from, or develop

subsequent to) infection with such microbes as *Staphylococcus aureus*, *Chlamydia trachomatis*, *Yersinia* e.g., *Y. enterocolitica*, *Salmonella*, e.g., *S. enteritidis*, *Shigella* and *Campylobacter*. *S. aureus* has been reported to be the major human pathogen in septic arthritis, responsible for the majority of cases.

- 5 IL-4 and IL-4-dependent Th2 responses play roles in promoting septic arthritis. IL-4 antagonist(s) are employed in accordance with the invention, to inhibit IL-4 and also to suppress the Th2 response in patients having septic arthritis or at risk for developing septic arthritis.

IL-4 increases bacterial burden and bacterial persistence in joints, by inhibiting clearance of the bacteria. IL-4 antagonists may be employed to assist in the clearance of bacteria associated  
10 with reactive arthritis, thereby reducing clinical manifestations such as swelling in joints. IL-4 antagonists may be administered to a human patient afflicted with septic arthritis, to reduce IL-4 mediated joint inflammation. In one approach, an antagonist is injected into a joint, e.g., into synovial fluid in the knee.

The use of IL-4 antagonists may benefit patients having (or at risk for) septic arthritis by  
15 suppressing a TH2 response and promoting a TH1 response against the infection. TH2 cytokines may contribute to bacterial persistence in the joint, whereas a TH1 response plays a role in eliminating the bacteria.

The antagonists may be administered to patients infected with bacteria or other microbes such as those listed above, to prevent development of septic arthritis. Antagonist(s) may be  
20 administered after diagnosis with such an infection, but before development of clinical symptoms of septic arthritis.

#### Whipple's Disease

*Tropheryma whippellii* is the causative bacterium for Whipple's Disease, also known as  
25 intestinal lipodystrophy and lipophagia granulomatosis. The disease is characterized by steatorrhea, frequently generalized lymphadenopathy, arthritis, fever, and cough. Also reported in Whipple's Disease patients are an abundance of "foamy" macrophages in the jejunal lamina propria, and lymph nodes containing periodic acid-schiff positive particles appearing bacilliform by electron microscopy (*Steadman's Medical Dictionary*, 26<sup>th</sup> Edition, Williams & Wilkins, Baltimore,  
30 MD, 1995).

The use of IL-4 antagonist(s) may benefit patients having (or at risk for developing) Whipple's Disease, by restoring a normal balance between the TH1 and TH2 components of the patient's immune response. Increased production of IL-4 (a TH2-type cytokine) and decreased  
35 levels of certain TH1-type cytokines have been associated with Whipple's Disease. TH2 cytokines may contribute to bacterial persistence, whereas a TH1 response plays a role in clearing the causative bacteria. IL-4 antagonists may be administered to patients infected with *T. whippellii*, whether or not the patient exhibits clinical symptoms of Whipple's Disease.

#### Dermatitis herpetiformis

Dermatitis herpetiformis, also known as Duhring's disease, is a chronic skin condition characterized by blistering skin lesions, cutaneous IgA deposits, and itching. Patients have an immunobullous skin disorder with an associated gluten sensitive enteropathy, which is mediated by a Th2 immune response. IL-4 antagonist(s) are administered in accordance with the present invention, to inhibit IL-4 and the Th2 response, thus promoting healing of current lesions and reducing or preventing the formation of blisters on the extensor body surfaces.

#### Hypertrophic scarring

In accordance with the present invention, IL-4 antagonist(s) are administered to patients who have, or are susceptible to developing, hypertrophic scarring. In one method provided herein, an IL-4 antagonist is administered to a burn patient. An immune response to burns and other injury is believed to play a role in the pathogenesis of hypertrophic scarring. Increased production of TH2-type cytokines, including IL-4, and reduced levels of certain TH1-type cytokines have been reported in burn patients who have hypertrophic scarring. The use of IL-4 antagonists may benefit patients having (or at risk for developing) hypertrophic scarring, by suppressing a TH2-type immune response.

#### Urticaria

Urticaria, especially chronic forms thereof such as chronic idiopathic urticaria (CIU), may be treated with an IL-4 antagonist in accordance with the present invention. CIU patients have higher serum levels of IL-4 than controls, and may have a predominantly TH2-type cytokine profile. Mast cells and Th2-type T cells are implicated as primary effector cells in chronic urticaria. IL-4 stimulates mast cell proliferation. Mast cell degranulation leads to histamine release, subsequent erythema, eosinophilia, redness of skin, and itching. IL-4 antagonists are administered to inhibit IL-4 and reduce the TH2-type response, thereby helping to control a patient's urticaria.

#### Ulcerative colitis; other disorders of the gastrointestinal tract

IL-4 is implicated in the pathogenesis of ulcerative colitis. TH2-type cytokines including IL-4 may predominate in the colonic mucosa of patients with this disorder. The use of IL-4 antagonist(s) to suppress the TH2 response may alleviate this condition.

In addition to ulcerative colitis, other disorders of the gastrointestinal tract or digestive system may be treated with IL-4 antagonist(s). Examples of such disorders include, but are not limited to, inflammatory bowel disease (IBD), with ulcerative colitis and Crohn's Disease being forms of IBD, gastritis, ulcers, and mucosal inflammation.

Any gastrointestinal condition in which IL-4 plays a role may be treated with an IL-4 antagonist in accordance with the present invention. For example, conditions involving IL-4-induced inflammation of part of the gastrointestinal tract may be treated with an IL-4 antagonist.

Particular embodiments are directed to treatment of chronic inflammatory conditions in the gastrointestinal tract.

Other embodiments are directed to conditions in which IL-4-induced barrier disruption plays a role, e.g., conditions characterized by decreased epithelial barrier function in at least a portion of the gastrointestinal tract. Such conditions may, for example, involve damage to the epithelium that is induced by IL-4, directly or indirectly.

The intestinal epithelium forms a relatively impermeable barrier between the lumen and the submucosa. Disruption of the epithelial barrier has been associated with conditions such as inflammatory bowel disease. See the discussion in Youakim, A. and M. Ahdieh (*Am. J. Physiol.* 276 (*Gastrointest. Liver Physiol.* 39): G1279-G1288, 1999), hereby incorporated by reference in its entirety. A damaged or "leaky" barrier can allow antigens to cross the barrier, which in turn elicits an immune response that may cause further damage to gastrointestinal tissue. Such an immune response may include recruitment of neutrophils or T cells, for example. An IL-4 antagonist may be administered to inhibit undesirable stimulation of an immune response.

#### Lung disorders

Methods for treating IL-4-induced pulmonary disorders are provided herein. Such disorders include, but are not limited to, lung fibrosis, including chronic fibrotic lung disease, other conditions characterized by IL-4-induced fibroblast proliferation or collagen accumulation in the lungs, pulmonary conditions in which a Th2-type immune response plays a role, conditions characterized by decreased barrier function in the lung (e.g., resulting from IL-4-induced damage to the epithelium), or conditions in which IL-4 plays a role in an inflammatory response.

Cystic fibrosis is characterized by the overproduction of mucus and development of chronic infections. Inhibiting IL-4 and the Th2 response will reduce mucus production and help control infections such as allergic bronchopulmonary aspergillosis (ABPA).

Allergic bronchopulmonary mycosis occurs primarily in patients with cystic fibrosis or asthma, where a Th2 immune response is dominant. Inhibiting IL-4 and the Th2 response will help clear and control these infections.

Chronic obstructive pulmonary disease is associated with mucus hypersecretion and fibrosis. Inhibiting IL-4 and the Th2 response will reduce the production of mucus and the development of fibrosis thereby improving respiratory function and delaying disease progression.

Bleomycin-induced pneumopathy and fibrosis, and radiation-induced pulmonary fibrosis are disorders characterized by fibrosis of the lung which is manifested by the influx of Th2, CD4<sup>+</sup> cells and macrophages, which produce IL-4 which in turn mediates the development of fibrosis. Inhibiting IL-4 and the Th2 response will reduce or prevent the development of these disorders.

Pulmonary alveolar proteinosis is characterized by the disruption of surfactant clearance. IL-4 increases surfactant product. Use of IL-4 antagonists will decrease surfactant production and decrease the need for whole lung lavage.



Adult respiratory distress syndrome (ARDS) may be attributable to a number of factors, one of which is exposure to toxic chemicals. One patient population susceptible to ARDS is critically ill patients who go on ventilators. ARDS is a frequent complication in such patients. IL-4 antagonists may alleviate ARDS by reducing inflammation and adhesion molecules, although methods for treating such patients in accordance with the present invention are not limited by a particular mechanism of action. IL-4 antagonists may be used to prevent or treat ARDS.

Sarcoidosis is characterized by granulomatus lesions. Use of IL-4 antagonists to treat sarcoidosis, particularly pulmonary sarcoidosis, is contemplated herein.

Conditions in which IL-4-induced barrier disruption plays a role (e.g., conditions characterized by decreased epithelial barrier function in the lung) may be treated with IL-4 antagonist(s). Damage to the epithelial barrier in the lungs may be induced by IL-4 directly or indirectly. The epithelium in the lung functions as a selective barrier that prevents contents of the lung lumen from entering the submucosa. A damaged or "leaky" barrier allows antigens to cross the barrier, which in turn elicits an immune response that may cause further damage to lung tissue. Such an immune response may include recruitment of eosinophils or mast cells, for example. An IL-4 antagonist may be administered to inhibit such undesirable stimulation of an immune response.

IL-4 antagonists may be employed to promote healing of lung epithelium, thus restoring barrier function. IL-4 antagonists may be employed to promote healing of lung epithelium in asthmatics, for example. Alternatively, the antagonist is administered for prophylactic purposes, to prevent IL-4-induced damage to lung epithelium.

#### Tuberculosis

A TH2-type immune response is implicated in playing a role in causing tissue damage (e.g., necrosis of lung tissue) in tuberculosis (TB) patients. Elevated levels of IL-4 are associated with TB. IL-4 production may be particularly elevated in cavitary tuberculosis (i.e., in TB patients who have developed pulmonary cavities, which can be detected/visualized by such techniques as radiographs of the chest).

IL-4 antagonists may benefit TB patients (especially those with cavitary TB) by suppressing a TH2-type immune response, or by binding (and inactivating) excess secreted IL-4. Methods for treating such patients in accordance with the present invention are not limited by a particular mechanism of action, however. IL-4 antagonists advantageously are administered in an amount that restores the desired balance between the TH1 and TH2 components of the immune response, and reduces IL-4-induced tissue damage in a patient.

#### Churg-Strauss syndrome

Churg-Strauss syndrome, a disease also known as allergic granulomatous angiitis, is characterized by inflammation of the blood vessels in persons with a history of asthma or allergy, and by eosinophilia. IL-4 antagonist(s) may be administered to alleviate inflammation in patients

with this syndrome. The use of IL-4 antagonists to suppress a TH2-type immune response, and to combat eosinophilia, would benefit the patients.

#### Pre-eclampsia

- 5 Pre-eclampsia is a toxemia of late pregnancy. The condition is characterized by a sharp rise in blood pressure, generally accompanied by edema and albuminuria, during the third term of pregnancy.

Elevated TH1-type and TH2-type immune responses may play a role in the condition. One method provided herein comprises administering an IL-4 antagonist to a pregnant woman  
10 who has developed pre-eclampsia. The IL-4 antagonist is administered in an amount, and for a period of time, sufficient to reduce the level of IL-4 (or of TH2-type cytokines collectively) to a level that is considered normal during pregnancy. In general, the IL-4 antagonist is administered repeatedly throughout the duration of the pregnancy.

#### Scleroderma

- 15 IL-4 antagonist(s) are administered to scleroderma patients in accordance with the invention. The antagonists reduce IL-4-induced collagen synthesis by fibroblasts in the patients. The antagonists may be employed in preventing or reducing fibrosis in skin and lung tissues, as well as other tissues in which fibrosis occurs in scleroderma patients, suppressing collagen  
20 synthesis in such tissues, and in treating scleroderma-related pulmonary disease.

#### Benign Prostate Hyperplasia

- Benign prostate hyperplasia (BPH), also known as benign prostate hypertrophy, may be treated with IL-4 antagonist(s). While not wishing to be bound by a particular mechanism of action,  
25 administration of an IL-4 inhibitor may benefit a patient with BPH by suppressing IL-4-induced inflammation, or by suppressing a TH2-type immune response.

#### Grave's Disease

- Antibodies directed against thyrotropin receptor play an important role in Grave's Disease,  
30 a disorder characterized by hyperthyroidism. Studies of cytokine production in Grave's Disease patients show a shift toward a TH2-type cytokine response. Use of an IL-4 antagonist to suppress the TH2-type immune response, and suppress antibody production, would benefit Grave's Disease patients.

#### Sickle Cell Disease

- 35 Sickle cell disease patients typically experience intermittent periods of acute exacerbation called crises, with the crises being categorized as anemic or vaso-occlusive. IL-4 antagonists find use in treating or preventing sickle cell crisis, especially in patients with elevated IL-4 levels or in whom the immune response has shifted toward a TH2-type response. Sickle cell disease

(especially sickle cell crisis) has been associated with increased susceptibility to infectious diseases, including bacterial infections. Administering IL-4 antagonists to sickle cell disease patients may help the patient mount an immune response against infectious diseases.

5                   Sjogren's syndrome

The autoimmune disease known as Sjogren's syndrome or sicca syndrome typically combines dry eyes and dry mouth with a disorder of the connective tissues, such as rheumatoid arthritis, lupus, scleroderma, or polymyositis. The vast majority of patients are middle age (or older) females. Sjogren's syndrome is an inflammatory disease of glands (e.g., lacrimal and salivary glands) and other tissues of the body. The syndrome typically is associated with autoantibody production.

10                   IL-4 antagonists may be administered to reduce the inflammatory response (such as inflammation of glands, including lacrimal glands) in such patients. IL-4 antagonists may benefit Sjogren's syndrome patients by suppressing a TH2-type immune response, or by binding (and inactivating) excess IL-4 at inflammatory lesions. Methods for treating patients in accordance with  
15                   the present invention are not limited by a particular mechanism of action, however.

Autoimmune lymphoproliferative syndrome

20                   Manifestations of autoimmune lymphoproliferative syndrome include lymphoproliferation and autoantibody production. Patients with the syndrome reportedly have an inherited deficiency in apoptosis. IL-4 antagonists may benefit patients with this syndrome by suppressing a TH2-type immune response, or by binding (and inactivating) excess IL-4 at sites of inflammation. Methods for treating such patients in accordance with the present invention are not limited by a particular mechanism of action, however.

25

Autoimmune hemolytic anemia

Excessive IL-4 secretion, and a deficiency in TH1-type cytokines, are implicated in contributing to the pathogenesis of autoimmune hemolytic anemia. IL-4 antagonists are administered in accordance with the present invention, to benefit the patients by reducing autoantibody production, and by restoring a more normal balance between the TH1 and TH2 components of the immune response.

30

Autoimmune uveitis

35                   Uveitis involves inflammation of the uvea (generally considered to include the iris, ciliary body, and choroid, considered together). Excess IL-4 secretion is implicated as playing a role in pathogenesis of this sight-threatening inflammatory eye disease. In accordance with the present invention, IL-4 antagonist(s) are administered to a uveitis patient to reduce disease severity. In one embodiment, IL-4 antagonist(s) are administered to an individual who has autoimmune uveoretinitis.

### Kawasaki Disease

Also known as the mucocutaneous lymph node syndrome, Kawasaki disease (KD) mainly afflicts young children. The disease is characterized by particular changes in the mucus membranes lining the lips and mouth, and by enlarged, tender lymph glands. Symptoms typically include fever, conjunctivitis, inflammation of the lips and mucous membranes of the mouth, swollen glands in the neck, and a rash covering the hands and feet, leading to hardened, swollen and peeling skin on hands and feet. In children with Kawasaki Disease (KD), inflammation of arteries (vasculitis) may develop. Due to the effect of the disease on the vascular system, KD reportedly is the main cause of acquired heart disease in children.

IL-4 antagonists may be administered to patients with Kawasaki Disease, to reduce the elevated levels of IL-4 in the patient. Excessive IL-4 secretion and a deficiency in TH1-type cytokines contribute to the pathogenesis of the disease.

### Barrett's esophagus

Barrett's esophagus is a condition characterized by alteration (subsequent to irritation) of the cells in the epithelial tissue that lines the lower portion of the esophagus. Frequent reflux of the stomach contents into the esophagus, over time, can lead to Barrett esophagus. Patients with Barrett esophagus are at risk for developing esophageal cancer (e.g., adenocarcinoma). While not wishing to be bound by a particular mechanism of action, administration of an IL-4 antagonist may benefit a patient with Barrett's esophagus by suppressing a TH2-type immune response. In one embodiment, an IL-4 antagonist is administered to a patient with esophagitis, to inhibit progression to Barrett's esophagus.

### Nephrosis

Nephrosis, also known as nephrotic syndrome, is kidney disease that is non-inflammatory and non-malignant. In the condition known as minimal change nephrosis, glomerular damage (believed to arise from structural changes in glomerular visceral epithelial cells) results in abnormalities that include proteinuria. A TH2-type immune response (especially secretion of the TH2-type cytokines IL-4 and IL-13) are implicated as playing a role in pathogenesis of minimal change nephrosis.

### Other Indications

Additional examples of conditions that may be treated in accordance with the present invention include but are not limited to the following. IL-4 antagonists may be employed in treating or preventing hyper IgE syndrome, idiopathic hypereosinophil syndrome, allergic reactions to medication, autoimmune blistering diseases (e.g., pemphigus vulgaris or bullous pemphigoid), myasthenia gravis (an autoimmune muscular disease), and chronic fatigue syndrome. IL-4 inhibitors may be employed in treating GVHD; particular methods for treating GVHD combination

therapy with other therapeutic agents as described below. IL-4 inhibitors also find use in treating or preventing hepatotoxicity induced by drugs such as diclofenac (a non-steroidal anti-inflammatory drug).

- An IL-4 antagonist may be employed as an adjuvant to allergy immunotherapy treatment.
- 5 IL-4 antagonists find further use as vaccine adjuvants, such as adjuvants for cancer vaccines and infectious disease vaccines. The use of IL-4 antagonists is especially advantageous when favoring a TH1-type immune response would be beneficial in preventing or treating the condition for which the vaccine is being administered. IL-4 antagonists may be employed when reducing an antibody-mediated immune response and/or promoting a T-cell-mediated immune response is
- 10 desired.

#### IL-4 Antagonists

- IL-4 antagonists that may be employed in accordance with the present invention include compounds that inhibit a biological activity of IL-4. The IL-4-induced biological activities to be
- 15 inhibited by the methods provided herein are activities that directly or indirectly play a role in the condition to be treated.

Examples of IL-4 antagonists include, but are not limited to, IL-4 receptors (IL-4R), antibodies, other IL-4-binding molecules, and IL-4 muteins as discussed further below. The antibodies may bind IL-4 or may bind an IL-4 receptor, for example.

- 20 Antagonists that bind IL-4 include but are not limited to IL-4 receptors and anti-IL-4 antibodies. Endogenous IL-4 that becomes bound to such an antagonist is thereby prevented from binding its natural receptor on cell surfaces *in vivo*, and thus cannot manifest IL-4-mediated biological activities.

- Different types of antagonists may act at different sites or by different mechanisms of
- 25 action. Examples include but are not limited to antagonists that interfere with binding of IL-4 to cell surface receptors or that inhibit signal transduction. The site of action may be intracellular (e.g., by interfering with an intracellular signaling cascade), on a cell surface, or extracellular. Antagonists that act by interfering with the interaction of IL-4 with IL-4R may bind to either IL-4 or to the receptor. An antagonist need not completely inhibit an IL-4 induced activity to find use in the
- 30 present invention; rather, antagonists that reduce a particular activity of IL-4 are contemplated for use as well.

- The above-presented discussions of particular mechanisms of action for IL-4 antagonists in treating particular diseases are illustrative only, and the methods presented herein are not bound thereby. The mechanisms of action by which IL-4 antagonists ameliorate diseases are not
- 35 limited to those discussed above.

In treating particular disorders, an IL-4 antagonist may reduce the amount of active IL-4 at a particular site within the body that is involved in the disorder. Antagonists that bind IL-4 such that it no longer can bind to endogenous cellular receptors functionally reduce the amount of active IL-4 available for inducing biological responses.

An IL-4 antagonist may alleviate a disorder by reducing the amount of free endogenous IL-4 that is circulating in the body, e.g., in the bloodstream or in a particular tissue. When the action of IL-4 on such tissue plays a role in pathogenesis of the disease, the antagonist serves to block action of IL-4 in the tissue, thereby alleviating the disorder. In a further example, antagonists may inhibit IL-4-induced recruitment of cells to a site or tissue within the body, wherein such recruitment plays a role in causing or exacerbating a disease. The antagonists may inhibit an IL-4-mediated influx of cells involved in an immune or inflammatory response. An antagonist may act by reducing proliferation, activation, migration, influx, or accumulation of a particular cell type, or by inhibiting a biological response directly or indirectly attributable to a particular cell type. Examples of particular cell types are fibroblasts, mast cells, and eosinophils.

As discussed above, some conditions may be treated by suppressing a TH2-type immune response. IL-4 is associated with a TH2 response, and is one of the cytokines secreted by T-helper cells of type 2 (TH2 cells). An IL-4 antagonist may be administered to reduce a TH2-type immune response. The IL-4 antagonist may be said to reduce proliferation of TH2 cells, to suppress a TH2 response, to shift the immune response toward a TH1 response, or to favor a TH1-type response. The use of antagonists of other cytokines associated with a TH2-type immune response is discussed below. Antagonists of other TH2-type cytokine(s), such as IL-5, IL-10, or IL-13, may be administered to patients who have a disorder involving elevated levels of such cytokines. Techniques for measuring the amount of such cytokines in a patient, e.g., in the patient's serum, are well known.

One embodiment of the invention is directed to a method for inhibiting IL-4-induced damage to epithelium, comprising administering an IL-4 antagonist to an individual who has, or is at risk of developing, a condition in which IL-4-mediated epithelial barrier disruption plays a role. In accordance with the present invention, barrier function studies revealed that IL-4 plays a role in reduction of barrier function in models of lung epithelium and intestinal epithelium, and that a soluble human IL-4 receptor polypeptide (an IL-4 antagonist) inhibits the IL-4-mediated reduction of barrier function (see example 7).

Particular embodiments of methods provided herein comprise administering an IL-4 antagonist to inhibit IL-4-induced damage to epithelium in the gastrointestinal tract or lung. Such methods may be employed to prevent epithelial damage, or to restore epithelial barrier function (i.e., promote repair or healing of the epithelium). The ability of an IL-4 antagonist to inhibit IL-4-induced damage to epithelium may be confirmed in any of a number of suitable assays, such as those described in example 7 below.

Any inflammation associated with (or subsequent to) an infection also may be treated with an IL-4 antagonist. The antagonist may be administered to inhibit any IL-4-induced component of an inflammatory response resulting from microbial infection in the gastrointestinal tract, for example.

Combinations of two or more antagonists may be employed in methods and compositions of the present invention. Examples of suitable IL-4 antagonists are as follows.

### IL-4 Receptor

A preferred IL-4 antagonist is an IL-4 receptor (IL-4R). When administered *in vivo*, IL-4R polypeptides circulate in the body and bind to circulating endogenous IL-4 molecules, preventing interaction of IL-4 with endogenous cell surface IL-4 receptors, thus inhibiting transduction of IL-4-induced biological signals.

IL-4 receptors are described in U.S. Patent 5,599,905; Idzerda et al., *J. Exp. Med.* 171:861-873, March 1990 (human IL-4R); and Mosley et al., *Cell* 59:335-348, October 20, 1989 (murine IL-4R); each of which is hereby incorporated by reference. The protein described in those three references is sometimes referred to in the scientific literature as IL-4R $\alpha$ . Unless otherwise specified, the terms "IL-4R" and "IL-4 receptor" as used herein encompass this protein in various forms that are capable of functioning as IL-4 antagonists, including but not limited to soluble fragments, fusion proteins, oligomers, and variants that are capable of binding IL-4, as described in more detail below.

The nucleotide sequence of a human IL-4R cDNA, and the amino acid sequence encoded thereby, are set forth in Figures 1A-1C. The cDNA clone was isolated from a cDNA library derived from a CD4<sup>+</sup>/CD8<sup>+</sup> human T cell clone designated T22, as described in Idzerda et al., *J. Exp. Med.*, 171:861, March 1990, and in U.S. Patent 5,599,905, which are hereby incorporated by reference in their entirety. The DNA and amino acid sequences of Figures 1A-1C are presented in SEQ ID NO:1 and SEQ ID NO:2, respectively.

The encoded human IL-4R protein comprises (from N- to C-terminus) an N-terminal signal peptide, followed by an extracellular domain, a transmembrane region, and a cytoplasmic domain. The transmembrane region, which is underlined in Figure 1A, corresponds to amino acids 208 through 231. The cytoplasmic domain comprises amino acids 232 through 800.

A signal peptide includes amino acids -25 to -1 of SEQ ID NO:2. An alternative signal peptide cleavage site occurs between residues -3 and -2 of SEQ ID NO:2, such that the signal peptide corresponds to residues -25 through -3.

As is recognized in the pertinent field, the signal peptide cleavage site for a given protein may vary according to such factors as the particular expression system (especially the host cells) in which the protein is expressed. The exact boundaries of the signal peptide, and thus the extracellular domain, of a given recombinant protein thus may depend on the expression system employed. Further, the signal peptide may be cleaved at more than one position, generating more than one species of polypeptide in a preparation of recombinant protein.

In one embodiment, in which an expression vector comprises DNA encoding amino acids -25 through 207 of SEQ ID NO:2, the expressed recombinant IL-4R includes two species of mature soluble human IL-4R. The expressed polypeptides include a major species corresponding to amino acids -2 to 207 and a minor species corresponding to amino acids 1 to 207 of SEQ ID NO:2. Two alternate forms of the extracellular domain of human IL-4R thus correspond to residues -2 to

207 and 1 to 207 of SEQ ID NO:2. The term "mature" refers to a protein in a form lacking a signal peptide or leader sequence, as is understood in the pertinent art.

Among the IL-4 receptors suitable for use herein are IL-4R fragments. Truncated IL-4R polypeptides may occur naturally, e.g., as a result of proteolytic cleavage, post-translational processing, or alternative splicing of mRNA. Alternatively, fragments may be constructed by deleting terminal or internal portions of an IL-4R sequence, e.g., via recombinant DNA technology. Fragments that retain the ability to bind IL-4 may be identified in conventional binding assays. Such fragments may be soluble fragments, as discussed below.

In a preferred embodiment of the invention, the antagonist comprises a soluble form of the IL-4R. A soluble IL-4 receptor is a polypeptide that is secreted from the cell in which it is expressed, rather than being retained on the cell surface. The full length human IL-4R protein of SEQ ID NO:2 is a transmembrane protein, which, as described above, comprises an N-terminal signal peptide, followed by an extracellular domain, a transmembrane region, and a C-terminal cytoplasmic domain. Soluble IL-4R polypeptides lack the transmembrane region that would cause retention on the cell, and the soluble polypeptides consequently are secreted into the culture medium. The transmembrane region and intracellular domain of IL-4R may be deleted or substituted with hydrophilic residues to facilitate secretion of the receptor into the cell culture medium.

Particular embodiments of soluble IL-4R polypeptides lack the transmembrane region but comprise the extracellular domain (the complete extracellular domain or a fragment thereof that is capable of binding IL-4). As one option, the polypeptide comprises all or part of the cytoplasmic domain, as well as the extracellular domain (or fragment of the extracellular domain), but lacks the transmembrane region.

Examples of soluble human IL-4R polypeptides include, but are not limited to, polypeptides comprising amino acid residues x to y of SEQ ID NO:2, wherein x represents 1 or -2 and y represents an integer from 197 to 207. Preferred embodiments include polypeptides comprising residues 1 to 207 or -2 to 207 of SEQ ID NO:2.

A protein preparation administered as an IL-4 antagonist may comprise more than one form of IL-4R. For example, the preparation may comprise polypeptide molecules consisting of amino acids 1 to 207 of SEQ ID NO:2, as well as polypeptides consisting of amino acids -2 to 207 of SEQ ID NO:2.

IL-4R polypeptides arising from alternative mRNA constructs, e.g., which can be attributed to different mRNA splicing events following transcription, and which yield polypeptide translates capable of binding IL-4, are among the IL-4R polypeptides disclosed herein. Such alternatively spliced mRNAs may give rise to soluble polypeptides.

Further examples of IL-4 receptors that may be employed in the methods provided herein are variants having amino acid sequences which are substantially similar to the native interleukin-4 receptor amino acid sequence of SEQ ID NO:2, or fragments thereof. Variant IL-4 receptor



polypeptides that are capable of functioning as IL-4 antagonists may be employed in the methods of the present invention.

Any of a number of conventional assay techniques may be employed to confirm that a given form of IL-4R (e.g., an IL-4R fragment or variant) functions as an IL-4 antagonist. Examples include binding assays or assays that test the ability of a given IL-4R polypeptide to inhibit transduction of an IL-4-induced biological signal. Examples of suitable *in vitro* assays are described below.

"Substantially similar" IL-4 receptors include those having amino acid or nucleic acid sequences that vary from a native sequence by one or more substitutions, deletions, or additions, but retain a desired biological activity of the IL-4R protein. Examples of nucleic acid molecules encoding IL-4 receptors include, but are not limited to: (a) DNA derived from the coding region of a native mammalian IL-4R gene; (b) DNA that is capable of hybridization to a DNA of (a) under moderately stringent conditions and which encodes an IL-4R having a biological activity of a native IL-4R; or (c) DNA that is degenerate as a result of the genetic code to a DNA defined in (a) or (b) and which encodes an IL-4R having a biological activity of a native IL-4R. Due to code degeneracy, there can be considerable variation in nucleotide sequences encoding the same amino acid sequence.

Variants may be naturally occurring, such as allelic variants or those arising from alternative splicing of mRNA. Alternatively, variants may be prepared by such well known techniques as *in vitro* mutagenesis.

A variant sequence identified by Idzerda et al., *supra*, comprises a GTC codon encoding the amino acid valine (Val) at position 50, instead of isoleucine (Ile). The variant sequence is otherwise identical to the sequence of SEQ ID NOS:1 and 2. IL-4R fragments, such as soluble fragments, comprising Val at position 50 are provided herein.

In particular embodiments, an IL-4 receptor DNA or amino acid sequence is at least 80 percent identical to the sequence of a native IL-4R. Preferably, an IL-4R DNA or polypeptide comprises a sequence that is at least 90 percent identical to a native IL-4R DNA or amino acid sequence. One example is a human IL-4R comprising an amino acid sequence that is at least 80 percent identical to the sequence presented in SEQ ID NO:2. Another example is a soluble IL-4R comprising an amino acid sequence at least 80 percent identical to the sequence of the extracellular domain of human IL-4R. Further examples are polypeptides comprising amino acid sequences that are at least 90 percent identical to the sequence presented in SEQ ID NO:2, or a fragment thereof. In a particular embodiment, the polypeptide comprises no more than 10 amino acid substitutions. IL-4R polypeptides that retain the ability to bind IL-4 may be identified in conventional binding assays.

Percent similarity or percent identity may be determined, for example, by comparing DNA or amino acid sequence information using the GAP computer program, version 6.0, available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith

and Waterman (*Adv. Appl. Math.* 2:482, 1981). Briefly, the GAP program defines similarity as the number of aligned symbols (i.e., nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, ed., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

IL-4R polypeptides that vary from native proteins but possess a desired property may be constructed by, for example, substituting or deleting residues not needed for the particular biological activity. Substitutions may be conservative substitutions, such that a desired biological property of the protein is retained. Amino acids may be replaced with residues having similar physicochemical characteristics.

Cysteine residues can be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Other alterations of a native sequence involve modification of adjacent dibasic amino acid residues, to enhance expression in yeast host cells in which KEX2 protease activity is present.

The present invention also includes IL-4R with or without associated native-pattern glycosylation. The glycosylation pattern may vary according to the type of host cells in which the protein is produced. Another option is inactivation of N-glycosylation sites by site-specific mutagenesis. N-glycosylation sites in eukaryotic proteins are characterized by the amino acid triplet Asn-A<sub>1</sub>-Z, where A<sub>1</sub> is any amino acid except Pro, and Z is Ser or Thr. In this sequence, asparagine provides a side chain amino group for covalent attachment of carbohydrate. Such a site can be eliminated by substituting another amino acid for Asn or for residue Z, deleting Asn or Z, or inserting a non-Z amino acid between A<sub>1</sub> and Z, or an amino acid other than Asn between Asn and A<sub>1</sub>.

Oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Examples of techniques for making such alterations are described in Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Patent Nos. 4,518,584 and 4,737,462.

IL-4 receptors that may be employed also include derivatives, e.g., various structural forms of the primary protein which retain a desired biological activity. Due to the presence of ionizable amino and carboxyl groups, for example, an IL-4R protein may be in the form of acidic or basic salts, or in neutral form. Individual amino acid residues may also be modified by oxidation or reduction. The primary amino acid structure may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups

and the like, or by creating amino acid sequence mutants. PEGylated derivatives (modified with polyethylene glycol) are contemplated. Covalent derivatives may be prepared by linking particular functional groups to IL-4R amino acid side chains or at the N- or C-termini. IL-4R derivatives may also be obtained by cross-linking agents, such as M-maleimidobenzoyl succinimide ester and N-hydroxysuccinimide, at cysteine and lysine residues. IL-4R proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromide-activated, bisoxirane-activated, carbonyldiimidazole-activated or tosyl-activated agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking).

Other derivatives of IL-4R within the scope of this invention include covalent or aggregative conjugates of IL-4R or its fragments with other proteins or polypeptides, such as by expression of recombinant fusion proteins comprising heterologous polypeptides fused to the N-terminus or C-terminus of an IL-4R polypeptide. For example, the conjugated peptide may be a heterologous signal (or leader) polypeptide, e.g., the yeast  $\alpha$ -factor leader, or a peptide such as an epitope tag. IL-4R-containing fusion proteins can comprise peptides added to facilitate purification or identification of IL-4R (e.g., poly-His). Specific examples of poly-His fusion constructs that is biologically active are soluble human IL-4R (e.g., comprising residues -2 to 207 or 1-207 of SEQ ID NO:2) His His and soluble human IL-4R His His His His His His. An amino acid sequence of IL-4 receptor can also be linked to the Flag<sup>®</sup> peptide Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (DYKDDDDK) (SEQ ID NO:3) as described in Hopp et al., *Bio/Technology* 6:1204, 1988, and U.S. Patent 5,011,912. The Flag<sup>®</sup> peptide is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. Reagents useful for preparing fusion proteins in which the Flag<sup>®</sup> peptide is fused to a given polypeptide are commercially available (Sigma, St. Louis, MO).

Oligomers that contain IL-4R polypeptides may be employed as IL-4 antagonists. Oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers. Oligomers comprising two or more IL-4R polypeptides are contemplated for use, with one example being a homodimer. Other oligomers include heterodimers, heterotrimers, and the like, which comprise an IL-4R polypeptide as well as at least one polypeptide that is not derived from the IL-4R of SEQ ID NO:2.

One embodiment is directed to oligomers comprising multiple IL-4R polypeptides joined via covalent or non-covalent interactions between peptide moieties fused to the IL-4R polypeptides. Such peptides may be peptide linkers (spacers), or peptides that have the property of promoting oligomerization. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of IL-4R polypeptides attached thereto, as described in more detail below.

In particular embodiments, the oligomers comprise from two to four IL-4R polypeptides. The IL-4R moieties of the oligomer may be in any of the forms described above, e.g., variants or fragments. Preferably, the oligomers comprise soluble IL-4R polypeptides.

As one alternative, an oligomer is prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al. (*PNAS USA* 88:10535, 1991); Byrn et al. (*Nature* 344:677, 1990); and Hollenbaugh and Aruffo ("Construction of Immunoglobulin Fusion Proteins", in *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1 - 10.19.11, 1992).

One embodiment of the present invention is directed to a dimer comprising two fusion proteins created by fusing IL-4R to the Fc region of an antibody. A gene fusion encoding the IL-4R/Fc fusion protein is inserted into an appropriate expression vector. IL-4R/Fc fusion proteins are expressed in host cells transformed with the recombinant expression vector, and allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield divalent IL-4R.

The term "Fc polypeptide" as used herein includes native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization are also included. Fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns.

One suitable Fc polypeptide, described in PCT application WO 93/10151 (hereby incorporated by reference), is a single chain polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Patent 5,457,035 and in Baum et al., (*EMBO J.* 13:3992-4001, 1994). The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

In other embodiments, IL-4R may be substituted for the variable portion of an antibody heavy or light chain. If fusion proteins are made with both heavy and light chains of an antibody, it is possible to form an oligomer with as many as four IL-4R extracellular regions.

Soluble recombinant fusion proteins comprising an IL-4R and various portions of the constant region of an immunoglobulin are described in EP 464,533, along with procedures for preparing such fusion proteins and dimers thereof. Among the fusion proteins described in EP 464,533 are those comprising the extracellular portion of human IL-4R and an Fc polypeptide.

Alternatively, the oligomer is a fusion protein comprising multiple IL-4R polypeptides, with or without peptide linkers (spacer peptides). Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233.

Another method for preparing oligomeric IL-4R involves use of a leucine zipper. Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., *Science* 240:1759, 1988), and have since been found in a variety of different proteins. Among the

known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe et al. (*FEBS Letters* 344:191, 1994), hereby  
5 incorporated by reference. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow et al. (*Semin. Immunol.* 6:267-278, 1994). In one approach, recombinant fusion proteins comprising a soluble IL-4R polypeptide fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric IL-4R that forms is recovered from the culture supernatant.

One example of a heterodimer comprises an IL-4R polypeptide derived from the human IL-4R of SEQ ID NO:2, and an IL-2R $\gamma$  polypeptide. IL-2R $\gamma$  (also known as IL-2R $\gamma$ ) is described in U.S. Patent 5,510,259 and in Takeshita et al. (*Science* 257:379, 17 July 1992), which are  
10 incorporated by reference herein. The polypeptides may be in one of the various forms described herein, e.g., soluble fragments, variants, and the like, derived from the indicated proteins. One embodiment of such a heterodimer comprises a soluble IL-4R/Fc fusion protein and a soluble IL-2R $\gamma$ /Fc fusion protein. Such heterodimers are described in WO 96/11213, along with IL-4R homodimers.

Other examples of heterodimers comprise an IL-4R subunit (preferably a soluble fragment of the protein of SEQ ID NO:2) and at least one IL-13 receptor subunit. IL-13 receptor (IL-13R)  
20 complexes and IL-13R polypeptides (such as polypeptides designated IL-13R $\alpha$ 1 and IL-13R $\alpha$ 2) are described in Zurawski et al., *J. Biol. Chem.* 270 (23), 13869, 1995; de Vries, *J. Allergy Clin. Immunol.* 102(2):165, August 1998; Caillard et al. *Immunology Today*, 17(3):108, March 1996, and U.S. Patent 5,710,023, each of which is incorporated by reference herein. In one embodiment, a heterodimer comprises a soluble human IL-4R and a soluble IL-13R (preferably a soluble form of  
25 the polypeptide described in U.S. Patent 5,710,023 or IL-13R $\alpha$ 1). The components of heterodimers may be any suitable form of the polypeptides that retains the desired activity, such as fragments, variants, or fusion proteins (e.g., fusions of soluble receptor polypeptides with Fc polypeptides, leucine zipper peptides, peptide linkers, or epitope tags).

IL-4 receptor polypeptides and fusion proteins described herein may be prepared by any  
30 of a number of conventional techniques. IL-4R polypeptides may be purified from cells that naturally express the receptor (such as the cells discussed in Park et al., *Proc. Natl. Acad. Sci. USA* 84:1669-673, 1987), or may be produced in recombinant expression systems, using well known techniques. Expression systems for use in producing IL-4R include those described in U.S. Patent 5,599,905, which is hereby incorporated by reference.

A variety of expression systems are known for use in producing recombinant proteins. In  
35 general, host cells are transformed with a recombinant expression vector that comprises DNA encoding a desired IL-4R polypeptide. Among the host cells that may be employed are prokaryotes, yeast or higher eukaryotic cells. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include insect cells and

established cell lines of mammalian origin. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, 293 cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, BHK (ATCC CRL 10) cell lines, and the CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10:2821, 1991). Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985).

The transformed cells are cultured under conditions that promote expression of the IL-4R, and the polypeptide is recovered by conventional protein purification procedures. One such purification procedure includes the use of affinity chromatography, e.g., over a matrix having IL-4 bound thereto. Expressed IL-4R will be deposited in the cell membrane or secreted into the culture supernatant, depending on the IL-4R DNA selected. Polypeptides contemplated for use herein include substantially homogeneous recombinant mammalian IL-4R polypeptides substantially free of contaminating endogenous materials

#### Antibodies

Antibodies that function as IL-4 antagonists may be employed in the methods of the present invention. The antibodies preferably are monoclonal antibodies or antigen-binding fragments thereof. Advantageously, humanized or chimeric monoclonal antibodies are employed. Most preferred are human monoclonal antibodies prepared using transgenic mice, as described below. Human antibodies also may be prepared using phage display techniques.

Examples of suitable antibodies are those that interfere with the binding of IL-4 to an IL-4 receptor. Such antibodies, referred to herein as blocking antibodies, may be raised against either IL-4 or IL-4R, and screened in conventional assays for the ability to interfere with binding of IL-4 to IL-4 receptors. Examples of suitable assays are assays that test the antibodies for the ability to inhibit binding of IL-4 to cells expressing IL-4R, or that test antibodies for the ability to reduce a biological or cellular response that results from the binding of IL-4 to cell surface IL-4 receptors.

It has been reported that IL-4R $\alpha$  is a component of certain multi-subunit IL-13 receptor complexes (Zurawski et al., *J. Biol. Chem.* 270 (23), 13869, 1995; de Vries, *J. Allergy Clin. Immunol.* 102(2):165, August 1998; and Callard et al. *Immunology Today*, 17(3):108, March 1996, each incorporated by reference herein). Thus, some antibodies raised against IL-4R $\alpha$  may interfere with the binding of IL-13 to such receptor complexes.

In one embodiment, antibodies directed against IL-4R block binding of IL-4 and also IL-13 to cells. The antibodies inhibit IL-4-induced biological activity and also inhibit IL-13-induced activity, and thus may be employed in treating conditions induced by either or both cytokines. Examples of such conditions include but are not limited to IgE-mediated conditions, asthma, allergic conditions, allergic rhinitis, and dermatitis including atopic dermatitis.

Antibodies that bind to IL-4R and inhibit IL-4 binding may be screened in various conventional assays to identify those antibodies that also interfere with the binding of IL-13 to such receptor complexes. Antibodies may be screened in binding assays or tested for the ability to inhibit an IL-4-induced and an IL-13-induced biological activity. An example of a suitable assay is illustrated in Example 5 below.

Antibodies specific for IL-4 or IL-4R may be prepared by well known procedures. See, for example, *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet et al. (eds.), Plenum Press, New York (1980); and *Antibodies: A Laboratory Manual*, Harlow and Land (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1988).

Antigen-binding fragments of such antibodies may be produced by conventional techniques. Examples of such fragments include, but are not limited to, Fab, F(ab')<sub>2</sub>, and Fv fragments. Antibody fragments and derivatives produced by genetic engineering techniques are also contemplated for use. Unless otherwise specified, the terms "antibody" and "monoclonal antibody" as used herein encompass both whole antibodies and antigen-binding fragments thereof.

Additional embodiments include chimeric antibodies, e.g., humanized versions of murine monoclonal antibodies. Such humanized antibodies may be prepared by known techniques, and offer the advantage of reduced immunogenicity when the antibodies are administered to humans. In one embodiment, a humanized monoclonal antibody comprises the variable region of a murine antibody (or just the antigen binding site thereof) and a constant region derived from a human antibody. Alternatively, a humanized antibody fragment may comprise the antigen binding site of a murine monoclonal antibody and a variable region fragment (lacking the antigen-binding site) derived from a human antibody. Procedures for the production of chimeric and further engineered monoclonal antibodies include those described in Riechmann et al. (*Nature* 332:323, 1988), Liu et al. (*PNAS* 84:3439, 1987), Larrick et al. (*BioTechnology* 7:934, 1989), and Winter and Harris (*TIPS* 14:139, May, 1993).

A method for producing an antibody comprises immunizing a non-human animal, such as a transgenic mouse, with an IL-4R polypeptide, whereby antibodies directed against the IL-4R polypeptide are generated in said animal. Procedures have been developed for generating human antibodies in non-human animals. The antibodies may be partially human, or preferably completely human. For example, transgenic mice into which genetic material encoding one or more human immunoglobulin chains has been introduced may be employed. Such mice may be genetically altered in a variety of ways. The genetic manipulation may result in human immunoglobulin polypeptide chains replacing endogenous immunoglobulin chains in at least some (preferably virtually all) antibodies produced by the animal upon immunization.

Mice in which one or more endogenous immunoglobulin genes have been inactivated by various means have been prepared. Human immunoglobulin genes have been introduced into the mice to replace the inactivated mouse genes. Antibodies produced in the animals incorporate

human immunoglobulin polypeptide chains encoded by the human genetic material introduced into the animal.

Examples of techniques for production and use of such transgenic animals are described in U.S. Patents 5,814,318, 5,569,825, and 5,545,806, which are incorporated by reference herein.

- 5 Examples 2-4 below provide further description of the preparation of transgenic mice useful for generating human antibodies directed against an antigen of interest.

Antibodies produced by immunizing transgenic non-human animals with an IL-4R polypeptide are provided herein. Transgenic mice into which genetic material encoding human immunoglobulin polypeptide chain(s) has been introduced are among the suitable transgenic  
10 animals. Examples of such mice include, but are not limited to, those containing the genetic alterations described in the examples below. One example of a suitable immunogen is a soluble human IL-4R, such as a polypeptide comprising the extracellular domain of the protein of SEQ ID NO:2, or other immunogenic fragment of the protein of SEQ ID NO:2.

Monoclonal antibodies may be produced by conventional procedures, e.g., by  
15 immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells may be fused with myeloma cells to produce hybridomas, by conventional procedures.

A method for producing a hybridoma cell line comprises immunizing such a transgenic animal with an IL-4R immunogen; harvesting spleen cells from the immunized animal; fusing the  
20 harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; and identifying a hybridoma cell line that produces a monoclonal antibody that binds an IL-4R polypeptide. Such hybridoma cell lines, and anti-IL-4R monoclonal antibodies produced therefrom, are encompassed by the present invention. Monoclonal antibodies secreted by the hybridoma cell line are purified by conventional techniques. Hybridomas or MAbs may be further screened to identify MAbs with  
25 particular properties, such as the ability to block an IL-4-induced activity, and to block an IL-13-induced activity (see the assay in example 5).

Human antibodies that bind IL-4R are provided by the present invention. In one embodiment of the invention, human antibodies raised against IL-4R and produced by techniques involving use of transgenic mice, block binding of IL-4 and also IL-13 to cells. Such antibodies are  
30 IL-4 antagonists and additionally function as IL-13 antagonists.

Among the uses of antibodies directed against an IL-4R is use in assays to detect the presence of IL-4R polypeptides, either *in vitro* or *in vivo*. The antibodies also may be employed in purifying IL-4R proteins by immunoaffinity chromatography. Those antibodies that additionally can block binding of IL-4 to IL-4R may be used to inhibit a biological activity that results from such  
35 binding. Blocking antibodies find use in the methods of the present invention. Such antibodies which function as IL-4 antagonists may be employed in treating any IL-4-induced condition, including but not limited to asthma and allergies, e.g., allergic rhinitis, contact dermatitis, and atopic dermatitis. In one embodiment, a human anti-IL-4R monoclonal antibody generated by procedures involving immunization of transgenic mice is employed in treating such conditions.



Antibodies may be employed in an *in vitro* procedure, or administered *in vivo* to inhibit an IL-4-induced biological activity. Disorders caused or exacerbated (directly or indirectly) by the interaction of IL-4 with cell surface IL-4 receptors thus may be treated. A therapeutic method involves *in vivo* administration of a blocking antibody to a mammal in an amount effective for

5 reducing an IL-4-induced biological activity.

Antibodies of the invention include, but are not limited to, partially human (preferably fully human) monoclonal antibodies that inhibit a biological activity of IL-4 and also inhibit a biological activity of IL-13. One embodiment is directed to a human monoclonal antibody that at least partially blocks binding of IL-4 to a cell, and at least partially blocks binding of IL-13 to a cell.

10 Antibodies of the present invention include but are not limited to antibodies generated by immunizing a transgenic mouse with an IL-4 receptor immunogen, wherein the transgenic mouse is selected from the mouse strains described in example 3 below. The desired antibodies are at least partially human, and preferably fully human. In one embodiment, the immunogen is a human IL-4 receptor polypeptide. Hybridoma cell lines derived from the thus-immunized mice, wherein

15 the hybridoma secretes a monoclonal antibody that binds IL-4R, also are provided herein. Examples of antibodies produced by immunizing such transgenic mice are the human monoclonal antibodies designated 6-2 (described in example 6); 12B5 (described in example 8); and MAbs 63, 1B7, 5A1, and 27A1 (all described in example 9). Monoclonal antibodies 6-2, 12B5, 63, 1B7, 5A1, and 27A1 are fully human antibodies, and are capable of inhibiting activity of both IL-4 and IL-13.

20 MAbs 12B5, 63, and 1B7 are preferred antagonists of human IL-4 and human IL-13.

Particular monoclonal antibodies of the invention are selected from the group consisting of MAb 6-2; a Mab that is cross-reactive with 6-2; a MAb that binds to the same epitope as 6-2; a MAb that competes with 6-2 for binding to a cell that expresses human IL-4R; a Mab that possesses a biological activity of 6-2; and an antigen-binding fragment of any of the foregoing

25 antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 6-2 for human IL-4R. MAb 6-2 is an IgM antibody. MAbs of other isotypes (including but not limited to IgG1 and IgG4), derived from 6-2, also are encompassed by the present invention. Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

30 One example of a biological activity of 6-2 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 6-2; and possesses IL-13-blocking activity substantially equivalent to that of 6-2. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

35 The DNA sequence of the variable region of the light chain of MAb 6-2 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable region of the heavy chain of MAb 6-2 is presented as SEQ ID NO:7, and the encoded amino acid sequence is presented in SEQ ID NO:8. Antibodies of the present invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain,

residues 1 to 107 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 118 of SEQ ID NO:8.

Complementarity determining regions (CDRs) of a given antibody may be identified using the system described by Kabat et al. in *Sequences of Proteins of Immunological Interest*, 5<sup>th</sup> Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication no. 91-3242, 1991). Particular embodiments of antibodies of the present invention comprise, within the variable region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 6-2. CDRs of 6-2 are discussed in example 6. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-35 of SEQ ID NO:6; residues 51-57 of SEQ ID NO:6; and residues 90-97 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 6-2. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-66; and residues 99-107 of SEQ ID NO:8.

Particular monoclonal antibodies of the invention are selected from the group consisting of MAb 12B5; a Mab that is cross-reactive with 12B5; a MAb that binds to the same epitope as 12B5; a MAb that competes with 12B5 for binding to a cell that expresses human IL-4R; a MAb that possesses a biological activity of 12B5; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 12B5 for human IL-4R. MAb 12B5 is an IgG1 antibody. MAbs of other isotypes, derived from 12B5, also are encompassed by the present invention. In particular embodiments, the isotype of the MAb is IgG1, IgG4, or IgM. Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

One example of a biological activity of 12B5 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 12B5, and possesses IL-13-blocking activity substantially equivalent to that of 12B5. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

IgG4 monoclonal antibodies derived from 12B5 are provided herein. Another embodiment is directed to IgM monoclonal antibodies derived from 12B5. Procedures for switching (altering) the subclass or isotype of an antibody are known in the pertinent field. Such procedures may involve, for example, recombinant DNA technology, whereby DNA encoding antibody polypeptide chains that confer the desired subclass is substituted for DNA encoding the corresponding polypeptide chain of the parent antibody.

The DNA sequence of the variable region of the light chain of MAb 12B5 is presented in SEQ ID NO:9, and the encoded amino acid sequence is presented in SEQ ID NO:10. The DNA sequence for the variable region of the heavy chain of MAb 12B5 is presented as SEQ ID NO:11, and the encoded amino acid sequence is presented in SEQ ID NO:12. Antibodies of the present

invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:10; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:12.

Particular embodiments of antibodies of the present invention comprise, within the variable region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 12B5. CDRs of 12B5 are discussed in example 8. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-35 of SEQ ID NO:10; residues 51-57 of SEQ ID NO:10; and residues 90-99 of SEQ ID NO:10. Particular antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 12B5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:12.

Particular monoclonal antibodies of the invention are selected from the group consisting of MAb 27A1; a Mab that is cross-reactive with 27A1; a Mab that binds to the same epitope as 27A1; a Mab that competes with 27A1 for binding to a cell that expresses human IL-4R; a Mab that possesses a biological activity of 27A1; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 27A1 for human IL-4R. 27A1 is an IgG1 antibody. MAbs of other isotypes, derived from 27A1, also are encompassed by the present invention. Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

One example of a biological activity of 27A1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 27A1; and possesses IL-13-blocking activity substantially equivalent to that of 27A1. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

The DNA sequence of the variable region of the light chain of MAb 27A1 is presented in SEQ ID NO:13, and the encoded amino acid sequence is presented in SEQ ID NO:14. The DNA sequence for the variable region of the heavy chain of MAb 27A1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:14; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 116 of SEQ ID NO:16.

Particular embodiments of antibodies of the present invention comprise, within the variable region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 27A1. CDRs of 27A1 are discussed in example 9. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-35 of SEQ ID

NO:14; residues 51-57 of SEQ ID NO:14; and residues 90-99 of SEQ ID NO:14. Particular antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 27A1. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-66; and residues 99-105 of SEQ ID NO:16.

Particular monoclonal antibodies of the invention are selected from the group consisting of MAb 5A1; a Mab that is cross-reactive with 5A1; a MAb that binds to the same epitope as 5A1; a MAb that competes with 5A1 for binding to a cell that expresses human IL-4R; a Mab that possesses a biological activity of 5A1; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 5A1 for human IL-4R. 5A1 is an IgG1 antibody. MAbs of other isotypes, derived from 5A1, also are encompassed by the present invention. Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

One example of a biological activity of 5A1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 5A1; and possesses IL-13-blocking activity substantially equivalent to that of 5A1. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

The DNA sequence of the variable region of the light chain of MAb 5A1 is presented in SEQ ID NO:17, and the encoded amino acid sequence is presented in SEQ ID NO:18. The DNA sequence for the variable region of the heavy chain of MAb 5A1 is presented as SEQ ID NO:19, and the encoded amino acid sequence is presented in SEQ ID NO:20. Antibodies of the present invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain, residues 1 to 107 of SEQ ID NO:18; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 123 of SEQ ID NO:20.

Particular embodiments of antibodies of the present invention comprise, within the variable region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 5A1. CDRs of 5A1 are discussed in example 9. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-34 of SEQ ID NO:18; residues 50-56 of SEQ ID NO:18; and residues 89-97 of SEQ ID NO:18. Particular antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 5A1. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-65; and residues 98-112 of SEQ ID NO:20.

Particular monoclonal antibodies of the invention are selected from the group consisting of MAb 63; a Mab that is cross-reactive with MAb 63; a MAb that binds to the same epitope as 63; a MAb that competes with 63 for binding to a cell that expresses human IL-4R; a Mab that

possesses a biological activity of 63; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 63 for human IL-4R. MAb 63 is an IgM antibody. MAbs of other isotypes, derived from 63, also are encompassed by the present invention.

5 Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

One example of a biological activity of 63 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 63; and possesses IL-13-blocking activity

10 substantially equivalent to that of 63. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

The DNA sequence of the variable region of the light chain of MAb 63 is presented in SEQ ID NO:21, and the encoded amino acid sequence is presented in SEQ ID NO:22. The DNA sequence for the variable region of the heavy chain of MAb 63 is presented as SEQ ID NO:23, and the encoded amino acid sequence is presented in SEQ ID NO:24. Antibodies of the present

15 invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain, residues 1 to 107 of SEQ ID NO:22; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 117 of SEQ ID NO:24.

Particular embodiments of antibodies of the present invention comprise, within the variable

20 region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 63. CDRs of 63 are discussed in example 9. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-34 of SEQ ID NO:22; residues 50-56 of SEQ ID NO:22; and residues 89-97 of SEQ ID NO:22. Particular

25 antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 63. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-66; and residues 99-106 of SEQ ID NO:24.

Particular monoclonal antibodies of the invention are selected from the group consisting of

30 MAb 1B7; a Mab that is cross-reactive with 1B7; a MAb that binds to the same epitope as 1B7; a MAb that competes with 1B7 for binding to a cell that expresses human IL-4R; a MAb that possesses a biological activity of 1B7; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of 1B7 for human IL-4R. MAbs of other isotypes,

35 derived from 1B7, also are encompassed by the present invention. Hybridoma cell lines that produce any such monoclonal antibodies also are provided by the present invention.

One example of a biological activity of 1B7 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, a MAb of the invention possesses IL-4-blocking activity substantially equivalent to that of 1B7; and possesses IL-13-blocking activity

substantially equivalent to that of 1B7. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in example 5).

MAb 1B7 was derived from MAb 63. The amino acid sequence of the heavy chain of MAb 1B7 is identical to that of MAb 63. The only differences between the two antibodies are in the light chain. The DNA sequence of the variable region of the light chain of MAb 1B7 is presented in SEQ ID NO:25, and the encoded amino acid sequence is presented in SEQ ID NO:26. The DNA sequence for the variable region of the heavy chain of MAb 1B7 is presented as SEQ ID NO:23, and the encoded amino acid sequence is presented in SEQ ID NO:24 (same as for MAb 63). Antibodies of the present invention include, but are not limited to, monoclonal antibodies that comprise, in their light chain, residues 1 to 107 of SEQ ID NO:26; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 117 of SEQ ID NO:24.

Particular embodiments of antibodies of the present invention comprise, within the variable region of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of 1B7. CDRs of 1B7 are discussed in example 9. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable region: amino acid residues 24-34 of SEQ ID NO:26; residues 50-56 of SEQ ID NO:26; and residues 89-97 of SEQ ID NO:26. Particular antibodies provided herein comprise, within the variable region of their heavy chain, at least one of the CDRs found in the heavy chain of 1B7. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable region: residues 31-35; residues 50-66; and residues 99-106 of SEQ ID NO:24.

Figure 4 presents an alignment of the amino acid sequences for the variable region of the light chain for antibodies 1B7, 63, 12B5, 6-2, 5A1, and 27A1. The first sequence in the alignment is for antibody 1B7. In the sequences for the other antibodies, amino acid residues that differ from the residue found at the corresponding position in the 1B7 sequence are shown, and residues that are identical to the residue found at that position in 1B7 are indicated by "-".

Figure 5 presents an alignment of the amino acid sequences for the variable region of the heavy chain for antibodies 1B7, 63, 12B5, 6-2, 5A1, and 27A1. The first sequence in the alignment is for antibody 1B7. In the sequences for the other antibodies, amino acid residues that differ from the residue found at the corresponding position in the 1B7 sequence are shown, and residues that are identical to the residue found at that position in 1B7 are indicated by "-".

As can be seen in Figures 4 and 5, a number of residues are highly conserved, in that the residue appears at a corresponding position in all six of the antibodies. Particular antibodies of the invention comprise, within the variable region of their light chain, amino acid residues that are found in the corresponding position in at least three of the sequences of Figure 4. In certain embodiments, the antibodies comprise residues found in a corresponding position in at least four, or at least five, or at least six, of the sequences of Figure 4. Particular antibodies of the invention comprise, within the variable region of their heavy chain, amino acid residues that are found in the corresponding position in at least three of the sequences of Figure 5. In certain embodiments, the

antibodies comprise residues found in a corresponding position in at least four, or at least five, or at least six, of the sequences of Figure 5.

Derivatives of monoclonal antibodies directed against IL-4R may be prepared, and screened for desired properties, by any of a number of known techniques. Certain of the techniques involve isolating DNA encoding a polypeptide chain (or portion thereof) of a MAb of interest, and manipulating the DNA through recombinant DNA technology. The DNA may be fused to another DNA of interest, or altered (e.g., by mutagenesis or other conventional techniques) to add, delete, or substitute one or more amino acid residues, for example.

DNA encoding antibody polypeptides (e.g., heavy or light chain, variable region only or full length) may be isolated from B-cells of mice that have been immunized with IL-4R. The DNA may be isolated by conventional procedures such as polymerase chain reaction (PCR). Phage display is another example of a known technique whereby derivatives of antibodies may be prepared. In one approach, polypeptides that are components of an antibody of interest are expressed in any suitable recombinant expression system, and the expressed polypeptides are allowed to assemble to form antibody molecules.

Single chain antibodies may be formed by linking heavy and light chain variable region (Fv region) fragments via an amino acid bridge (short peptide linker), resulting in a single polypeptide chain. Such single-chain Fvs (scFvs) have been prepared by fusing DNA encoding a peptide linker between DNAs encoding the two variable region polypeptides ( $V_L$  and  $V_H$ ). The resulting antibody fragments can form dimers or trimers, depending on the length of a flexible linker between the two variable domains (Kortt et al., *Protein Engineering* 10:423, 1997). Techniques developed for the production of single chain antibodies include those described in U.S. Patent No. 4,946,778; Bird (*Science* 242:423, 1988); Huston et al. (*Proc. Natl. Acad. Sci. USA* 85:5879, 1988); and Ward et al. (*Nature* 334:544, 1989). Single chain antibodies derived from antibodies provided herein (including but not limited to scFvs derived from MAbs 6-2, 12B5, 63, 1B7, 5A1, and 27A1) are encompassed by the present invention.

Techniques are known for deriving an antibody of a different subclass or isotype from an antibody of interest, i.e., subclass switching. Thus, IgG1 or IgG4 monoclonal antibodies may be derived from an IgM monoclonal antibody, for example, and vice versa. Such techniques allow the preparation of new antibodies that possess the antigen-binding properties of a given antibody (the parent antibody), but also exhibit biological properties associated with an antibody isotype or subclass different from that of the parent antibody. Recombinant DNA techniques may be employed. Cloned DNA encoding particular antibody polypeptides may be employed in such procedures, e.g., DNA encoding the constant region of an antibody of the desired isotype.

In particular embodiments, antibodies raised against IL-4R have a binding affinity ( $K_a$ ) for IL-4R of at least  $1 \times 10^8$ . In other embodiments, the antibodies exhibit a  $K_a$  of at least  $1 \times 10^8$  or at least  $1 \times 10^{10}$ .

PEGylated derivatives of antibodies (modified with polyethylene glycol) also are contemplated, and may be prepared by conventional techniques. Also provided herein are

conjugates comprising a detectable (e.g., diagnostic) or therapeutic agent, attached to an antibody directed against IL-4R. Examples of such agents are well known, and include but are not limited to diagnostic radionuclides, therapeutic radionuclides, and cytotoxic drugs. The conjugates find use in *in vitro* or *in vivo* procedures.

5 Particular embodiments of the invention are directed to novel nucleic acid molecules and polypeptides. DNA and amino acid sequence information has been determined for polypeptides that are components of certain antibodies of the present invention, as discussed in examples 6, 8, and 9 below. Among the nucleic acids of the present invention is isolated DNA comprising a nucleotide sequence selected from the group consisting of the nucleotide sequence presented in  
10 SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, and SEQ ID NO:25. Among the polypeptides of the present invention is a purified polypeptide comprising an amino acid sequence selected from the group consisting of the amino acid sequence presented in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID  
15 NO:20, SEQ ID NO:22, SEQ ID NO:24, and SEQ ID NO:26. For *in vivo* use, the polypeptides advantageously are purified. A polypeptide may be purified individually, or in the form of a purified antibody of which the polypeptide is a component.

Further examples of IL-4 antagonists are antibodies that bind IL-4 and inhibit the binding of IL-4 to cell surface receptors. Such antibodies may be prepared, and screened to identify those  
20 that are blocking antibodies, by conventional procedures. Antigen-binding fragments of such antibodies find use as antagonists, as do humanized or genetically engineered derivatives thereof.

Examples of procedures for preparing antibodies directed against human IL-4 (including monoclonal antibodies), assays by which blocking antibodies are identified, and techniques for generating humanized or genetically engineered derivatives of anti-IL-4 antibodies, are described  
25 in U.S. Patents 5,041,381, 5,863,537, 5,928,904, and 5,676,940, which are hereby incorporated by reference. Further examples of antibodies that may be employed as IL-4 antagonists are described in WO 91/09059, also incorporated by reference herein.

#### Other antagonists

30 Any compound that functions as an IL-4 antagonist and is suitable for administration in accordance with the methods of the present invention may be employed. Antagonists need not completely abolish IL-4-induced biological activity to be useful. Rather, a given antagonist may reduce a biological activity of IL-4.

Derivatives, mutants/mutins, and other variants of IL-4 that function as IL-4 antagonists  
35 may be employed. Peptides (which may or may not be mutins) derived from IL-4 that bind to an IL-4R without inducing transduction of a biological signal find use herein. Such peptides function as inert blockers, interfering with the binding of biologically active endogenous IL-4 to cell surface receptors. IL-4-induced signal transduction thereby is inhibited. Mutins or other antagonists that



induce a biological response at a reduced level or to a lesser degree, compared to the response induced by native IL-4, also find use as IL-4 antagonists.

In particular embodiments, a mutein of human IL-4 is capable of inhibiting IL-4-induced and IL-13-induced biological activity. Such muteins may be employed in those methods described below that involve use of an antagonist of both IL-4 and IL-13 to treat particular disorders. See, for example, the human IL-4 mutein (a variant polypeptide with two point mutations) described in Fitch et al. (*J. Allergy Clin. Immunol.*, Vol. 107, no. 2, pg. S61, abstract no. 209, 2001).

Further examples of IL-4 antagonists, including IL-4 muteins, and procedures for preparation thereof are described in Muller et al., *J. Mol. Biol.*, 237:423-436, 1994; U.S. Patent 6,028,176, and U.S. Patent 5,723,118, which are each incorporated by reference herein.

Other options are antisense molecules (oligonucleotides) that inhibit expression of IL-4. Alternatively, the antisense molecule may suppress expression of other molecules involved in IL-4-induced signal transduction.

Any suitable assay, including *in vitro* assays, can be utilized to determine whether a given compound inhibits an IL-4-induced biological activity. An antagonist may be assayed for the ability to inhibit <sup>3</sup>H-thymidine incorporation in cells that normally proliferate in response to IL-4.

An alternative involves use of conventional binding assay techniques to test an antagonist for the ability to inhibit binding of IL-4 to cells expressing native or recombinant IL-4 receptors. For use in such assays, recombinant human IL-4 can be expressed and purified as described in U.S. Patent 5,017,691, hereby incorporated by reference herein, or in Park et al., *J. Exp. Med.* 166:476 (1987). The purified protein may be labeled with a detectable agent (e.g., radiolabeled) by any of a number of conventional techniques. A commercially available enzymebead radioiodination reagent (BioRad) may be employed in radiolabeling IL-4 with <sup>125</sup>I for example.

The ability of an IL-4 antagonist to inhibit IL-4-induced damage to epithelium, such as lung epithelium or intestinal epithelium (which may result in loss of barrier function), may be confirmed in any of a number of suitable assays. Among the suitable assay techniques are those described in example 7 below.

#### Therapeutic Methods and Administration of Antagonists

Methods provided herein comprise administering an IL-4 antagonist to a patient, thereby reducing an IL-4-induced biological response that plays a role in a particular condition. In particular embodiments, methods of the invention involve contacting endogenous IL-4 with an IL-4 antagonist, e.g., in an *ex vivo* procedure.

Treatment encompasses alleviation of at least one symptom of a disorder, or reduction of disease severity, and the like. An antagonist need not effect a complete "cure", or eradicate every symptom or manifestation of a disease, to constitute a viable therapeutic agent. As is recognized in the pertinent field, drugs employed as therapeutic agents may reduce the severity of a given disease state, but need not abolish every manifestation of the disease to be regarded as useful therapeutic agents. One embodiment of the invention is directed to a method comprising

administering to a patient an IL-4 antagonist in an amount and for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of the particular disorder.

Antibodies that inhibit the binding of both IL-4 and IL-13 to cells are discussed herein. A method for suppressing IL-4-induced and IL-13-induced activities in humans comprises administering an effective amount of such an antibody. Conditions induced by IL-4 or by IL-13, or by both cytokines, thus may be treated.

As is understood in the pertinent field, antagonists are administered to a patient in a manner appropriate to the indication. Antagonists may be administered by any suitable technique, including but not limited to parenterally, topically, or by inhalation. If injected, the antagonist can be administered, for example, via intra-articular, intravenous, intramuscular, intralesional, intraperitoneal or subcutaneous routes, by bolus injection, or continuous infusion. Localized administration, e.g. at a site of disease or injury is contemplated, as are transdermal delivery and sustained release from implants. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation of the antagonist in aerosol form, and the like. Other alternatives include eyedrops; oral preparations including pills, syrups, lozenges or chewing gum; and topical preparations such as lotions, gels, sprays, and ointments.

Use of IL-4 antagonists in *ex vivo* procedures is contemplated. For example, a patient's blood (bodily fluid containing IL-4) may be contacted with an antagonist that binds IL-4 *ex vivo*, thereby reducing the amount of IL-4 in the fluid when returned to the patient. The antagonist may be bound to a suitable insoluble matrix or solid support material.

Advantageously, antagonists are administered in the form of a composition comprising at least one IL-4 antagonist and one or more additional components such as a physiologically acceptable carrier, excipient or diluent. The present invention provides such compositions comprising an effective amount of an IL-4 antagonist, for use in the methods provided herein.

The compositions contain antagonist(s) in any of the forms described herein. The antagonist may be a whole antibody or an antigen-binding fragment or engineered derivative thereof, for example. For compositions containing an IL-4 receptor, the receptor may be any of the fragments, variants, or oligomers of the protein of SEQ ID NO:2 described herein, for example.

Compositions may, for example, comprise an antagonist together with a buffer, antioxidant such as ascorbic acid, low molecular weight polypeptide (such as those having fewer than 10 amino acids), protein, amino acid, carbohydrate such as glucose, sucrose or dextrins, chelating agents such as EDTA, glutathione, and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are examples of appropriate diluents. In accordance with appropriate industry standards, preservatives such as benzyl alcohol may also be added. The composition may be formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Suitable components are nontoxic to recipients at the dosages and concentrations employed. Further examples of components that may be employed in

pharmaceutical formulations are presented in *Remington's Pharmaceutical Sciences*, 16<sup>th</sup> Ed., Mack Publishing Company, Easton, PA, 1980.

Kits for use by medical practitioners include an IL-4 antagonist and a label or other instructions for use in treating any of the conditions discussed herein. The kit preferably includes a sterile preparation of one or more IL-4 antagonists, which may be in the form of a composition as disclosed above, and may be in one or more vials.

Dosages and the frequency of administration may vary according to such factors as the route of administration, the particular antagonist employed, the nature and severity of the disease to be treated, whether the condition is acute or chronic, and the size and general condition of the patient. Appropriate dosages can be determined by procedures known in the pertinent art, e.g. in clinical trials that may involve dose escalation studies.

An antagonist may be administered once, or repeatedly. In particular embodiments, the antagonist is administered over a period of at least a month or more, e.g., for one, two, or three months or even indefinitely. For treating chronic conditions, long-term treatment is generally most effective. However, for treating acute conditions, administration for shorter periods, e.g. from one to six weeks, may be sufficient. In general, the antagonist is administered until the patient manifests a medically relevant degree of improvement over baseline for the chosen indicator or indicators.

Particular embodiments of the present invention involve administering an antagonist at a dosage of from about 1 ng/kg/day to about 10 mg/kg/day, more preferably from about 500 ng/kg/day to about 5 mg/kg/day, and most preferably from about 5 ug/kg/day to about 2 mg/kg/day, to a patient. In additional embodiments, an antagonist such as a soluble human IL-4R polypeptide is administered to adults one time per week, two times per week, or three or more times per week, to treat the medical disorders disclosed herein. If injected, the effective amount of antagonist per adult dose may range from 1-20 mg/m<sup>2</sup>, and preferably is about 5-12 mg/m<sup>2</sup>. Alternatively, a flat dose may be administered; the amount may range from 5-100 mg/dose. One range for a flat dose is about 20-30 mg per dose. In one embodiment of the invention, a flat dose of 25 mg/dose is repeatedly administered by injection. If a route of administration other than injection is used, the dose is appropriately adjusted in accordance with standard medical practices. One example of a therapeutic regimen involves injecting a dose of about 20-30 mg of IL-4R or other antagonist one to three times per week over a period of at least three weeks, though treatment for longer periods may be necessary to induce the desired degree of improvement. For pediatric patients (age 4-17), one suitable regimen involves the subcutaneous injection of 0.4 mg/kg, up to a maximum dose of 25 mg of IL-4R, administered two or three times per week.

Particular embodiments of the methods provided herein involve subcutaneous injection of from 0.5 mg to 10 mg, preferably from 3 to 5 mg, of a soluble IL-4R, once or twice per week. Another embodiment is directed to pulmonary administration (e.g., by nebulizer) of 3 or more mg of a soluble IL-4R once a week.

Examples of therapeutic regimens provided herein comprise subcutaneous injection of a soluble human IL-4R once a week, at a dose of 1.5 to 3 mg, to treat pulmonary sarcoidosis, minimal change nephrosis, autoimmune uveitis, sickle cell crisis, Churg-Strauss syndrome, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, pre-eclampsia, autoimmune hemolytic anemia, Barrett's esophagus, Grave's Disease, Kawasaki Disease, and cavitary tuberculosis. Weekly administration of IL-4R is continued until symptoms subside. Treatment may resume as needed, or, alternatively, maintenance doses may be administered.

An antagonist is administered to the patient in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the patient's illness may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. In most instances, an improvement is considered to be sustained if the patient exhibits the improvement on at least two occasions separated by two to four weeks. The degree of improvement generally is determined by the patient's physician, who may make this determination based on signs or symptoms, and who may also employ questionnaires that are administered to the patient, such as quality-of-life questionnaires developed for a given disease.

As one example, in treating benign prostate hyperplasia, an IL-4 inhibitor is administered to the patient in an amount and for a time effective in scar regression or complete healing. Maintenance doses may be given or treatment resumed as needed.

Elevated levels of IL-4 are associated with a number of disorders, as discussed above. Patients with a given disorder may be screened, to identify those individuals who have elevated IL-4 levels, or to identify those with an elevated TH2-type immune response, thereby identifying the patients who may benefit most from treatment with an IL-4 antagonist. Thus, treatment methods provided herein optionally comprise a first step of measuring a patient's IL-4 level. An IL-4 antagonist may be administered to a patient in whom IL-4 levels are elevated above normal. Alternatively or additionally, a patient may be pre-screened to determine whether the patient has an elevated TH2-type immune response, prior to administration of antagonist(s) against one or more TH2-type cytokines.

A patient's levels of IL-4 (and, optionally, of other TH2-type cytokines) may be monitored during and/or after treatment with an IL-4 antagonist, to detect reduction in the levels of the cytokines. For some disorders, the incidence of elevated IL-4 levels, and the balance between TH1-type and TH2-type immune responses, may vary according to such factors as the stage of the disease or the particular form of the disease. Known techniques may be employed for measuring IL-4 levels, e.g., in a patient's serum, and for assessing TH2-type immune responses. Cytokine levels in blood samples may be measured by ELISA, for example.

Particular embodiments of methods and compositions of the invention involve the use of two or more different IL-4 antagonists. In further embodiments, IL-4 antagonist(s) are

administered alone or in combination with other agents useful for treating the condition with which the patient is afflicted. Examples of such agents include both proteinaceous and non-proteinaceous drugs. When multiple therapeutics are co-administered, dosages may be adjusted accordingly, as is recognized in the pertinent art. "Co-administration" and combination therapy are not limited to simultaneous administration, but include treatment regimens in which an IL-4 antagonist is administered at least once during a course of treatment that involves administering at least one other therapeutic agent to the patient.

Examples of other agents that may be co-administered with IL-4 antagonists are other antibodies, cytokines, or cytokine receptors, which are chosen according to the particular condition to be treated. Alternatively, non-proteinaceous drugs that are useful in treating one of the particular conditions discussed above may be co-administered with an IL-4 antagonist.

For treating IgE-mediated conditions, an IL-4 antagonist may be co-administered with an IgE antagonist. One example is an anti-IgE antibody. Humanized anti-IgE monoclonal antibodies are described in Presta et al. (*J. Immunol.* 151(5):2623-2632, 1993) and Adelroth et al. (*J. Allergy Clin. Immunol.* 106(2):253-259, 2000), for example.

IL-4 antagonists may be co-administered with an IL-5 antagonist, which may be a molecule that interferes with the binding of IL-5 to an IL-5 receptor, such as an anti-IL-5 antibody (e.g., a human or humanized anti-IL-5 monoclonal antibody), or a receptor such as a soluble human IL-5 receptor polypeptide. IL-5 has been implicated as playing a role in mediating allergic responses. Thus, administration of antagonist(s) of IL-4 and IL-5 is contemplated for treatment of allergic reactions, including but not limited to allergic asthma.

IL-4 antagonists may be employed in conjunction with other agent(s) in treating the particular IL-4-induced conditions discussed above. For example, drugs currently employed in treating the conditions may be co-administered with one or more IL-4 antagonists.

For treating asthma, an IL-4 antagonist may be co-administered with other anti-asthma medications, such as inhaled corticosteroids, beta agonists, leukotriene antagonists, xanthines, fluticasone, salmeterol, albuterol, non-steroidal agents such as cromolyn, and the like. IL-4 antagonists may be co-administered with other anti-allergy medications to treat allergic reactions.

One embodiment of the present invention is directed to co-administration of an IL-4 antagonist (such as a soluble human IL-4R) and fluticasone and salmeterol to treat a disorder such as asthma. Compositions comprising an IL-4 inhibitor (e.g., soluble human IL-4R), fluticasone, and salmeterol are provided herein. Advair Diskus (Glaxo Wellcome) comprises fluticasone propionate and salmeterol xinafoate. For treating asthma, Advair Diskus and the IL-4 antagonist preferably are delivered by inhalation.

Another example of combination therapy comprises co-administration of an IL-4 antagonist and an IL-9 antagonist to a patient who has asthma. Any suitable IL-9 antagonist may be employed, such as an IL-9 receptor (preferably a soluble form thereof), an antibody that interferes with binding of IL-9 to a cell surface receptor (wherein the antibody may be raised against IL-9 or an IL-9 receptor polypeptide), or another compound that inhibits IL-9-induced biological activity.

IL-9 receptors include those described in WO 93/18047 and U.S. Patents 5,789,237 and 5,962,269, which are hereby incorporated by reference herein.

In an additional embodiment of combination therapy, a method for treating ulcerative colitis comprises co-administration of at least one IL-4 antagonist and at least one IL-1 antagonist. Examples of IL-1 antagonists include type I IL-1 receptor, type II IL-1 receptor, IL-1 receptor antagonist (IL-1Ra), antagonistic (blocking) antibodies directed against IL-1, and antagonistic antibodies directed against an IL-1 receptor. Various forms of the receptors may be employed, such as fragments, variants and fusions analogous to those described above for IL-4 receptor. A preferred IL-1 antagonist is a soluble form of type II IL-1 receptor, which is described in U.S. Patent 5,350,683, hereby incorporated by reference herein.

One method of the present invention comprises co-administering IL-4 antagonist(s) and IL-13 antagonist(s) to a patient who has minimal change nephrosis. Alternative embodiments involve administering IL-4 antagonist(s) alone, or IL-13 antagonist(s) alone, to a minimal change nephrosis patient. The IL-4 antagonist(s) and/or IL-13 antagonist(s) may be administered to reduce severity of the disease.

Another method provided herein is a method for treating various allergic inflammatory conditions, comprising co-administering IL-4 antagonist(s) and IL-13 antagonist(s). Conditions such as asthma, allergies, and chronic lung diseases such as cystic fibrosis and chronic obstructive pulmonary disease are treated by such a method.

Any suitable IL-13 antagonist may be employed, including but not limited to IL-13 receptors (preferably soluble forms thereof), IL-13 receptor antagonists, antibodies directed against IL-13 or an IL-13 receptor, other proteins that interfere with the binding of IL-13 to an IL-13 receptor, and compounds that inhibit IL-13-mediated signal transduction. IL-13 receptors and heterodimers comprising IL-13R polypeptides as components thereof are described above. Antibodies that are raised against IL-4R may be screened for the ability to also function as IL-13 antagonists, as discussed above.

A method for treating or preventing a condition characterized by reduced epithelial barrier function comprises co-administering IL-4 antagonist(s) and one or more IL-13 antagonists. Such conditions are discussed above. In one embodiment, the condition is asthma. Particular embodiments are directed to co-administering one or more IL-4 antagonists and one or more IL-13 antagonists to a patient having a condition involving reduction of lung epithelial barrier function or intestinal epithelial barrier function, wherein both IL-4 and IL-13 play a role in the reduced barrier function. The method thus inhibits both IL-4-induced reduction of barrier function and IL-13-induced reduction of barrier function. The adverse effect of IL-13 on lung and intestinal epithelial barrier function can be confirmed using assay techniques such as those described in example 7 below. (See also Zund et al., *J. Biol. Chem.* 271(13):7460-7464, 1996.)

Another method provided herein comprises co-administering IL-4 antagonist(s) and interferon- $\gamma$  (IFN- $\gamma$ ) to a patient having a condition involving reduction of lung epithelial barrier function. Optionally, such a method further comprises co-administering one or more IL-13

antagonists to the patient (i.e., co-administering an IL-4 antagonist, IFN- $\gamma$ , and an IL-13 antagonist). Other methods comprise administering IFN- $\gamma$  as a single agent, or co-administering IFN- $\gamma$  and an IL-13 antagonist, to a patient having a condition involving reduction of lung epithelial barrier function. In one embodiment, the patient has asthma. For treating asthma, the IL-4 antagonist, IFN- $\gamma$ , and/or IL-13 antagonist preferably are administered by inhalation.

One method provided herein for treating asthma comprises administering an IL-4 antagonist and interferon- $\gamma$  to a human who has asthma. Another method for treating asthma comprises co-administering an IL-4 antagonist, IFN- $\gamma$ , and an IL-13 antagonist to a human who has asthma. In one embodiment, IFN- $\gamma$  is co-administered to an asthmatic, together with an antibody that functions as an antagonist of both IL-4 and IL-13. Such antibodies are described elsewhere herein.

A single agent may function as an IL-4 antagonist and an IL-13 antagonist, as discussed above. As an example of such an agent, some antibodies raised against IL-4R $\alpha$  may interfere with the binding of both IL-4 and IL-13 receptor complexes, due to the shared IL-4R $\alpha$  component in such multi-subunit receptor complexes (discussed above). Thus, a single agent may be employed in a method for inhibiting reduction of barrier function.

Antagonists may be co-administered with one or more leukotriene receptor antagonists to treat disorders such as allergic inflammatory diseases, e.g., asthma and allergies. Examples of leukotriene receptor antagonists include but are not limited to montelukast, pranlukast, and zafirlukast. Drugs that function as 5-lipoxygenase inhibitors may be co-administered with an IL-4 antagonist to treat asthma.

Methods provided herein comprise administering one or more of the following to Churg-Strauss Syndrome patients: IL-4 antagonist(s), IL-5 antagonist(s), IL-13 antagonist(s) or IgE antagonist(s). One example of such a method involves co-administering IL-4 antagonist(s) and IL-5 antagonist(s) to a Churg-Strauss Syndrome patient. In another embodiment, IL-4 antagonist(s) and IgE antagonist(s) are co-administered to the patient. In yet another embodiment, IL-4 antagonist(s) and IL-13 antagonist(s) are co-administered to the patient.

The hormone relaxin may be co-administered with an IL-4 antagonist to treat scleroderma (systemic sclerosis), idiopathic pulmonary fibrosis, or any other disorder characterized by pulmonary fibrosis, such as the conditions involving fibrosis of the lung that are discussed above. Recombinant human relaxin is preferred for use in treating humans.

A method for treating benign prostate hyperplasia comprises co-administering IL-4 antagonist(s) and one or more additional anti-inflammatory agents. Examples of agents that inhibit inflammation include tumor necrosis factor (TNF) antagonists and IL-17 antagonists.

Any suitable IL-17 antagonist may be employed, including but not limited to an IL-17 receptor (preferably soluble forms thereof), IL-17 receptor antagonists, antibodies directed against IL-17 or an IL-17 receptor, other proteins that interfere with the binding of IL-17 to an IL-17 receptor, and compounds that inhibit IL-17-mediated signal transduction. An IL-17 receptor, including soluble forms thereof and oligomers thereof, is described in WO 96/29408, hereby

incorporated by reference. An alternative method provided herein comprises administering an IL-17 antagonist to treat a patient with benign prostate hyperplasia.

Likewise, any suitable TNF antagonist may be employed, including but not limited to a TNF receptor (preferably soluble forms thereof), fusion proteins comprising a TNF receptor (or comprising the TNF-binding portion of a TNF receptor), TNF receptor antagonists, antibodies directed against TNF or a TNF receptor, other proteins that interfere with the binding of TNF to a TNF receptor, and compounds that inhibit TNF-mediated signal transduction. Further examples of TNF inhibitors are the drugs thalidomide and pentoxifylline. The TNF receptor protein known as p75 or p80 TNF-R preferably is employed. A preferred TNF antagonist is a soluble human TNF receptor (sTNF-R) in dimeric form, such as dimers of sTNF-R/Fc fusion proteins. One such dimer is etanercept (Enbrel<sup>®</sup>, Immunex Corporation, Seattle, WA). p75/p80 TNF-R, including soluble fragments and other forms thereof, is described in WO 91/03553, hereby incorporated by reference herein.

In accordance with the present invention, an IL-4 antagonist is co-administered with a TNF antagonist to treat any condition in which undesirable IL-4-induced and TNF-induced immune responses play a role, such as inflammation. One method provided herein comprises co-administering an IL-4 antagonist and a TNF antagonist to a patient with inflammatory bowel disease, Crohn's disease, or ulcerative colitis. Other embodiments are directed to a method comprising co-administering an IL-4 antagonist and a TNF antagonist to a patient who has Kawasaki Disease, autoimmune hemolytic anemia, autoimmune uveoretinitis, autoimmune lymphoproliferative syndrome, Sjogren's syndrome, chronic fatigue syndrome, or hepatotoxicity induced by a drug such as diclofenac.

Another method provided herein comprises co-administering an IL-4 antagonist and a TNF antagonist to a pregnant woman who has developed pre-eclampsia. Administration of the IL-4 antagonist and TNF-antagonist preferably continues for the duration of the pregnancy.

Suitable dosages of etanercept (Enbrel<sup>®</sup>, Immunex Corporation, Seattle, WA) will vary according to the nature of the disease to be treated, disease severity, the size of the patient (e.g., adult or child), and other factors, as is recognized in the pertinent field. In one embodiment of the methods provided herein, Enbrel<sup>®</sup> is administered twice a week by subcutaneous injection at a dose of from 1 to 25 mg. One embodiment of a pediatric dosage is 0.4 mg/kg. Particular methods provided herein comprise co-administration of an IL-4 antagonist and Enbrel<sup>®</sup> to a patient has autoimmune lymphoproliferative syndrome or Sjogren's syndrome, wherein Enbrel<sup>®</sup> is given by subcutaneous injection at a dose of from 1 to 25 mg.

For treating graft versus host disease, an IL-4 antagonist is co-administered with at least one of the following agents: a TNF antagonist, an IL-1 antagonist, steroids, or corticosteroids. The TNF inhibitor preferably is Enbrel<sup>®</sup>. A preferred IL-1 antagonist is a soluble form of type II IL-1 receptor, which is described in U.S. Patent 5,350,683. In one embodiment, the GVHD is associated with (e.g., develops subsequent to) bone marrow transplantation. An IL-4 antagonist



may be employed in combination with at least one of the above-listed agents, in methods for suppressing an immune response directed against transplanted cells, tissue, and/or alloantigen.

A number of cytokine antagonists and other agents/drugs are disclosed herein as being useful for combination therapy (e.g., co-administration with an IL-4 antagonist) in treating particular diseases. It is to be understood that such antagonists, agents, or drugs also find use as single agents in treating those diseases. It also is to be understood that disclosure of methods involving administration of an antagonist to a particular cytokine, to treat a disease, encompasses administration of one type of antagonist, and also encompasses administration of two or more different antagonists for that cytokine, unless specified otherwise.

The following examples are offered by way of illustration, and not by way of limitation.

#### **EXAMPLE 1: Preparation of Monoclonal Antibodies**

IL-4 receptor polypeptides may be employed as immunogens in generating monoclonal antibodies by conventional techniques, e.g., techniques described in U.S. Patent 5,599,905, hereby incorporated by reference. It is recognized that polypeptides in various forms may be employed as immunogens, e.g., full length proteins, fragments thereof, fusion proteins thereof such as Fc fusions, cells expressing the recombinant protein on the cell surface, etc.

To summarize an example of such a procedure, an IL-4R immunogen emulsified in complete Freund's adjuvant is injected subcutaneously into Lewis rats, in amounts ranging from 10-100  $\mu$ l. Three weeks later, the immunized animals are boosted with additional immunogen emulsified in incomplete Freund's adjuvant and boosted every three weeks thereafter. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay, ELISA (enzyme-linked immunosorbent assay), or inhibition of binding of  $^{125}$ I-IL-4 to extracts of IL-4R-expressing cells. Following detection of an appropriate antibody titer, positive animals were given a final intravenous injection of antigen in saline. Three to four days later, the animals are sacrificed, splenocytes harvested, and fused to the murine myeloma cell line AG8653. The resulting hybridoma cell lines are plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

Hybridoma clones thus generated are screened for reactivity with IL-4R. Initial screening of hybridoma supernatants utilizes an antibody capture and binding of partially purified  $^{125}$ I-IL-4 receptor. Hybridomas that are positive in this screening method are tested by a modified antibody capture to detect hybridoma cells lines that are producing blocking antibody. Hybridomas that secrete a monoclonal antibody capable of inhibiting  $^{125}$ I-IL-4 binding to cells expressing IL-4R are thus detected. Such hybridomas then are injected into the peritoneal cavities of nude mice to produce ascites containing high concentrations (>1 mg/ml) of anti-IL-4R monoclonal antibody. The resulting monoclonal antibodies may be purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein G.

**EXAMPLE 2: Generation of Cmu targeted mice**

This example describes procedures for generating transgenic mice. Additional procedures for generating transgenic mice, and the use of such mice for preparing human antibodies, are described in Examples 3 and 4.

Construction of a CMD targeting vector. The plasmid pCEmu contains an EcoRI/XhoI fragment of the murine Ig heavy chain locus, spanning the mu gene, that was obtained from a Balb/C genomic lambda phage library (Marcu et al. *Cell* 22: 187, 1980). This genomic fragment was subcloned into the XhoI/EcoRI sites of the plasmid pCEmu9H (Marsh et al; *Gene* 32, 481-485, 1984). The heavy chain sequences included in pCEmu extend downstream of the EcoRI site located just 3' of the mu intronic enhancer, to the XhoI site located approximately 1 kb downstream of the last transmembrane exon of the mu gene; however, much of the mu switch repeat region has been deleted by passage in *E. coli*.

The targeting vector was constructed as follows. (See Figures 2A-2C, which depict further details.) A 1.3 kb HindIII/SmaI fragment was excised from pCEmu and subcloned into HindIII/SmaI digested pBluescript (Stratagene, La Jolla, CA). This pCEmu fragment extends from the HindIII site located approximately 1 kb 5' of Cmu1 to the SmaI site located within Cmu1. The resulting plasmid was digested with SmaI/SpeI and the approximately 4 kb SmaI/XbaI fragment from pCEmu, extending from the SmaI site in Cmu1 3' to the XbaI site located just downstream of the last Cmu exon, was inserted. The resulting plasmid, pTAR1, was linearized at the SmaI site, and a neo expression cassette inserted. This cassette consists of the neo gene under the transcriptional control of the mouse phosphoglycerate kinase (pgk) promoter (XbaI/TaqI fragment; Adra et al. (1987) *Gene* 60: 65-74) and containing the pgk polyadenylation site (PvuII/HindIII fragment; Boer et al. (1990) *Biochemical Genetics* 28: 299-308). This cassette was obtained from the plasmid pKJ1 (described by Tybulewicz et al. (1991) *Cell* 65: 1153-1163) from which the neo cassette was excised as an EcoRI/HindIII fragment and subcloned into EcoRI/HindIII digested pGEM-7Zf (+) to generate pGEM-7 (KJ1). The neo cassette was excised from pGEM-7 (KJ1) by EcoRI/SalI digestion, blunt ended and subcloned into the SmaI site of the plasmid pTAR1, in the opposite orientation of the genomic Cmu sequences.

The resulting plasmid was linearized with Not I, and a herpes simplex virus thymidine kinase (tk) cassette was inserted to allow for enrichment of ES clones bearing homologous recombinants, as described by Mansour et al. (1988) *Nature* 336: 348-352. This cassette consists of the coding sequences of the tk gene bracketed by the mouse pgk promoter and polyadenylation site, as described by Tybulewicz et al. (1991) *Cell* 65:1153-1163.

The resulting CMD targeting vector contains a total of approximately 5.3 kb of homology to the heavy chain locus and is designed to generate a mutant mu gene into which has been inserted a neo expression cassette in the unique SmaI site of the first Cmu exon. The targeting vector was linearized with PvuI, which cuts within plasmid sequences, prior to electroporation into ES cells.

Generation and analysis of targeted ES cells. AB-1 ES cells (McMahon, A. P. and Bradley, A., (1990) *Cell* 62: 1073-1085) were grown on mitotically inactive SNL76/7 cell feeder

layers (ibid.), essentially as described in *Teratocarcinomas and Embryonic Stem Cells: a Practical Approach*, E. J. Robertson, Ed., Oxford: IRL Press, 1987, pp. 71-112. The linearized CMD targeting vector was electroporated into AB-1 cells by the methods described in Hasty *et al.* (1991) *Nature* 350: 243-246. Electroporated cells were plated into 100 mm dishes at a density of  $1-2 \times 10^6$  cells/dish. After 24 hours, G418 (200 micrograms/ml of active component) and FIAU ( $5 \times 10^{-7}$  M) were added to the medium, and drug-resistant clones were allowed to develop over 8-9 days. Clones were picked, trypsinized, divided into two portions, and further expanded. Half of the cells derived from each clone were then frozen and the other half analyzed for homologous recombination between vector and target sequences.

DNA analysis was carried out by Southern blot hybridization. DNA was isolated from the clones as described by Laird *et al.*, (1991) *Nucleic Acids Res.* 19:4293). Isolated genomic DNA was digested with SpeI and probed with a 915 bp SacI fragment, probe A (Figure 2C), which hybridizes to a sequence between the mu intronic enhancer and the mu switch region. Probe A detects a 9.9 kb SpeI fragment from the wild type locus, and a diagnostic 7.6 kb band from a mu locus which has homologously recombined with the CMD targeting vector (the neo expression cassette contains a SpeI site).

Of 1132 G418 and FIAU resistant clones screened by Southern blot analysis, 3 displayed the 7.6 kb Spe I band indicative of homologous recombination at the mu locus. These 3 clones were further digested with the enzymes BglI, BstXI, and EcoRI to verify that the vector integrated homologously into the mu gene. When hybridized with probe A, Southern blots of wild type DNA digested with BglI, BstXI, or EcoRI produce fragments of 15.7, 7.3, and 12.5 kb, respectively, whereas the presence of a targeted mu allele is indicated by fragments of 7.7, 6.6, and 14.3 kb, respectively. All 3 positive clones detected by the SpeI digest showed the expected BglI, BstXI, and EcoRI restriction fragments diagnostic of insertion of the neo cassette into the Cmu1 exon.

Generation of mice bearing the mutated mu gene. The three targeted ES clones, designated number 264, 272, and 408, were thawed and injected into C57BL/6J blastocysts as described by A. Bradley in *Teratocarcinomas and Embryonic Stem Cells: a Practical Approach*, E. J. Robertson, Ed., Oxford: IRL Press, 1987, pp. 113-151. Injected blastocysts were transferred into the uteri of pseudopregnant females to generate chimeric mice representing a mixture of cells derived from the input ES cells and the host blastocyst. The extent of ES cell contribution to the chimera can be visually estimated by the amount of agouti coat coloration, derived from the ES cell line, on the black C57BL/6J background. Clones 272 and 408 produced only low percentage chimeras (i.e. low percentage of agouti pigmentation) but clone 264 produced high percentage male chimeras. These chimeras were bred with C57BL/6J females and agouti offspring were generated, indicative of germline transmission of the ES cell genome. Screening for the targeted mu gene was carried out by Southern blot analysis of BglI digested DNA from tail biopsies (as described above for analysis of ES cell DNA). Approximately 50% of the agouti offspring showed a hybridizing BglI band of 7.7 kb in addition to the wild type band of 15.7 kb, demonstrating a germline transmission of the targeted mu gene.

Analysis of transgenic mice for functional inactivation of mu gene. To determine whether the insertion of the neo cassette into Cmu1 has inactivated the Ig heavy chain gene, a clone 264 chimera was bred with a mouse homozygous for the JHD mutation, which inactivates heavy chain expression as a result of deletion of the JH gene segments (Chen et al, (1993) *Immunol.* 5: 647-656). Four agouti offspring were generated. Serum was obtained from these animals at the age of 1 month and assayed by ELISA for the presence of murine IgM. Two of the four offspring were completely lacking IgM (Table 1). Genotyping of the four animals by Southern blot analysis of DNA from tail biopsies by BglI digestion and hybridization with probe A (Figure 2C), and by Stul digestion and hybridization with a 475 bp EcoRI/Stul fragment (ibid.) demonstrated that the animals which fail to express serum IgM are those in which one allele of the heavy chain locus carries the JHD mutation, the other allele the Cmu1 mutation. Mice heterozygous for the JHD mutation display wild type levels of serum Ig. These data demonstrate that the Cmu1 mutation inactivates expression of the mu gene.

Table 1 presents the level of serum IgM, detected by ELISA, for mice carrying both the CMD and JHD mutations (CMD/JHD), for mice heterozygous for the JHD mutation (+JHD), for wild type (129Sv x C57BL/6J)F1 mice (+/+), and for B cell deficient mice homozygous for the JHD mutation (JHD/JHD).

TABLE 1

Mouse	Serum IgM (micrograms/ml)	Ig H chain genotype
42	<0.002	CMD/JHD
43	196	+JHD
44	<0.002	CMD/JHD
45	174	+JHD
129 x BL6 F1	153	+/+
JHD	<0.002	JHD/JHD

### EXAMPLE 3: Generation of transgenic mice

The HCo12 human heavy chain transgene. The HCo12 transgene was generated by coinjection of the 80 kb insert of pHc2 (Taylor et al., 1994, *Int. Immunol.*, 6: 579-591) and the 25 kb insert of pVx6. The plasmid pVx6 was constructed as described below.

An 8.5 kb HindIII/SalI DNA fragment, comprising the germline human VH1-18 (DP-14) gene together with approximately 2.5 kb of 5' flanking, and 5 kb of 3' flanking genomic sequence was subcloned into the plasmid vector pSP72 (Promega, Madison, WI) to generate the plasmid p343.7.16. A 7 kb BamHI/HindIII DNA fragment, comprising the germline human VH5-51 (DP-73) gene together with approximately 5 kb of 5' flanking and 1 kb of 3' flanking genomic sequence, was cloned into the pBR322 based plasmid cloning vector pGP1f (Taylor et al. 1992, *Nucleic Acids Res.* 20: 6287-6295), to generate the plasmid p251f.

A new cloning vector derived from pGP1f, pGP1k (the sequence of which is presented in Figures 3A and 3B and SEQ ID NO:4), was digested with EcoRV/BamHI, and ligated to a 10 kb EcoRV/BamHI DNA fragment, comprising the germline human VH3-23 (DP47) gene together with approximately 4 kb of 5' flanking and 5 kb of 3' flanking genomic sequence. The resulting plasmid, p112.2RR.7, was digested with BamHI/Sall and ligated with the 7 kb purified BamHI/Sall insert of p251f. The resulting plasmid, pVx4, was digested with XhoI and ligated with the 8.5 kb XhoI/Sall insert of p343.7.16. A clone was obtained with the VH1-18 gene in the same orientation as the other two V genes. This clone, designated pVx6, was then digested with NotI and the purified 26 kb insert collected, together with the purified 80 kb NotI insert of pH2 at a 1:1 molar ratio, into the pronuclei of one-half day (C57BL/6J x DBA/2J)F2 embryos as described by Hogan *et al.* (B. Hogan *et al.*, Manipulating the Mouse Embryo, A Laboratory Manual, 2<sup>nd</sup> edition, 1994, Cold Spring Harbor Laboratory Press, Plainview NY).

Three independent lines of transgenic mice comprising sequences from both Vx6 and HC2 were established from mice that developed from the injected embryos. These lines are designated (HCo12)14881, (HCo12)15083, and (HCo12)15087. Each of the three lines were then bred with mice comprising the CMD mutation described in Example 2, the JKD mutation (Chen *et al.* 1993, EMBO J. 12: 811-820), and the (KCo5)9272 transgene (Fishwild *et al.* 1996, Nature Biotechnology 14: 845-851). The resulting mice express human heavy and kappa light chain transgenes in a background homozygous for disruption of the endogenous mouse heavy and kappa light chain loci.

Additional transgenic mouse strains Particular strains of mice that may be used to generate IL-4R-reactive monoclonal antibodies are strain ((CMD)++; (JKD)++; (HCo7)11952+/+; (KCo5)9272+/+), and strain ((CMD)++; (JKD)++; (HCo12)15087+/+; (KCo5)9272+/+). Each of these transgenic strains is homozygous for disruptions of the endogenous heavy chain (CMD) and kappa light chain (JKD) loci. Both strains also comprise a human kappa light chain transgene (HCo7), with individual animals either hemizygous or homozygous for insertion #11952. The two strains differ in the human heavy chain transgene used. Mice were hemizygous or homozygous for either the HCo7 or the HCo12 transgene. The CMD mutation is described above in Example 2. The generation of (HCo12)15087 mice is described above (in this example). The JKD mutation (Chen *et al.* 1993, EMBO J. 12: 811-820) and the (KCo5)9272 (Fishwild *et al.* 1996, Nature Biotechnology 14: 845-851) and (HCo7)11952 mice, are described in U.S. Patent 5,770,429, which is hereby incorporated by reference.

#### EXAMPLE 4: Generation of Human Anti-IL-4R Monoclonal Antibodies

Transgenic mice Strain ((CMD)++; (JKD)++; (HCo7)11952+/+; (KCo5)9272+/+ which is homozygous for disruptions of the endogenous heavy chain (CMD) and kappa light chain (JKD) loci (see example 3), was used to generate IL-4R-reactive monoclonal antibodies. This strain also comprises a human kappa light chain transgene (HCo7) with individual animals either hemizygous

or homozygous for insertion #11952. Mice were hemizygous or homozygous for the HCo7 transgene. The CMD mutation is described above in Example 2. The JKD mutation (Chen *et al.* 1993, EMBO J. 12: 811-820) and the (KCo5)9272 (Fishwild *et al.* 1996, Nature Biotechnology 14: 845-851) and (HCo7)11952 mice, are described in U.S. Patent 5,770,429, which is hereby incorporated by reference.

Immunization. Transgenic mice were initially immunized i.p. with 25 ug IL-4R protein in adjuvant (Titermax, available from CytRx Corporation, Norcross, GA). The immunogen was a human IL-4R polypeptide comprising the extracellular domain of the protein of SEQ ID NO:2. Immunized mice were subsequently boosted every 4 weeks i.p. with the IL-4R immunogen in incomplete Freund's adjuvant. Animals were kept on protocol for 2 to 5 months. Prior to fusion, animals were boosted i.v. on days -4 and -3 with 5 to 8 ug immunogen.

Fusions. Spleen cells harvested from the immunized mice were fused to mouse myeloma cells NS-1 by standard procedures (Harlow and Lane, 1988, Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor New York; Kennett *et al.* 1980, Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analysis. Plenum, New York; Oi and Hertenberg, 1980, Immunoglobulin Producing Hybrid Cell Lines, in Selected Methods in Cellular Immunology, ed. Mishell and Shilgi, pp. 357-372. Freeman, San Francisco). Cells were cultured in DMEM, 10% FBS, OPI (Sigma O-5003), BME (Gibco 21985-023), 3% Origen Hybridoma Cloning Factor (Igen IG50-0615), and 5% P388d1 (ATCC TIB 63) conditioned media. HAT or HT supplement was added to the medium during initial growth and selection.

Hybridoma Screening. To identify hybridomas secreting human antibodies against the IL-4R, ELISA plates (Nunc MaxiSorp) were coated overnight at 4°C with 100 ul/well human IL-4R at 2.0 ug/ml in PBS. Plates were washed with 100 ul/well PBS-Tween (PBST) containing 1% BSA. Fifty ul cell culture supernatant was added followed by a 1.0 hour incubation. Plates were washed and then incubated for one hour with 100 ul/well goat anti-human IgG conjugated to horseradish peroxidase (Sigma #A-3813, or #A-7164). Plates were washed three times in PBS-Tween between each step.

Wells that read positive by ELISA were screened for their ability to block the binding of IL-4 to IL-4R. ELISA plates were coated overnight with a non-neutralizing mouse anti-human IL-4R antibody M10 at 2 ug/ml. Plates were washed 3X with PBST. 100 ul human IL-4R was added at 10 ng/ml in PBST and incubated for 1.0 hour. Plates were washed 4X with PBST and 100 ul supernatant samples were added and incubated for 1.0 hour. Wells were washed 4X with PBST. 5.0 ng/ml biotinylated IL-4 was added in PBST and incubated for 1.0 hour. 100 ul/well poly80 horseradish peroxidase (RDI) was added at 1:5000 in PBST and incubated for 45 minutes. Plates were washed 5X with PBST, and a colorimetric reagent (3,3',5,5' tetramethylbenzidine, available from Kirkegaard and Perry) was added at 100 ul/well until color developed. Reaction was stopped with 100 ul phosphoric acid and plates were read at 450nm. Absent or reduced signal was interpreted as the antibody binding to receptor in a manner that blocked IL-4 from

binding to receptor. Wells that appeared to block binding were expanded and tested for IL-4 and IL-13 blocking in a CD23 expression assay (see example 5).

#### EXAMPLE 5: Assay for assessing blocking activity

This assay is based on ability of both IL-4 and IL-13 to enhance the expression of the activation-associated surface antigen CD23 on human B cells. Antibodies are tested for the ability to inhibit CD23 expression induced by IL-4 and by IL-13.

Antibodies raised against human IL-4R (huIL-4R) were tested either in the form of hybridoma supernatants or purified protein. Prior to addition to cultures, the antibodies were buffer exchanged against culture medium (RPMI 1640 plus 10% heat-inactivated fetal bovine serum) by centrifugation, using Centricon filter devices (Amicon) with a 10kDa cutoff.

Human peripheral blood B cells were purified as described previously (Morris et al., *J. Biol. Chem.* 274:418-423, 1999). The B cells ( $3 \times 10^5$ /well) in culture medium were placed in 96-well round-bottomed microtiter plates and preincubated at room temperature for 30 min with test antibodies at the final concentrations indicated. Recombinant human IL-4 or IL-13 was then added to the cultures at the concentrations indicated, and cells were cultured for 20-24 hours at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>. At the end of the culture period, cells were washed once in PBS + 0.02% NaN<sub>3</sub> in the 96-well culture plate and were resuspended in blocking buffer (2% normal rabbit serum + 1% normal goat serum in PBS + NaN<sub>3</sub>). Phycoerythrin (PE)-conjugated CD23 monoclonal antibody (mAb) or PE-conjugated isotype control mAb (both from Pharmingen) was then added to cells at a final dilution of 1:10. Cells were incubated for 30 minutes at 4°C, washed x3 in PBS + NaN<sub>3</sub> and analyzed on a FacScan (Becton Dickinson) for CD23 expression.

In all experiments, negative controls were included which consisted of cells cultured with hybridoma growth medium or isotype-matched non-blocking human anti-hIL-4R antibody. An anti-huIL-4R murine mAb (R&D Systems), previously shown to block the binding and function of both hIL-4 and hIL-13, was used as a positive control for neutralization of CD23 induction by IL-4 and IL-13.

#### EXAMPLE 6: Hybridoma Cell Line

One hybridoma cell line generated by procedures described above (see example 4) is designated 6-2. The anti-IL-4R monoclonal antibody secreted by this hybridoma is a blocking antibody, as determined in a conventional plate binding assay, and thus functions as an IL-4 antagonist. The monoclonal antibody produced by 6-2 also exhibits the ability to reduce an IL-13-induced biological activity.

One embodiment of the invention is directed to a hybridoma cell line produced as described above, wherein the hybridoma secretes an isotype IgM MAb directed against human IL-4R. Also provided herein are IgG1 monoclonal antibodies derived from IgM monoclonal antibodies.

The DNA sequence of the variable region of the light chain of MAb 6-2 has been determined, and is presented in SEQ ID NO:5; the amino acid sequence encoded thereby is presented in SEQ ID NO:6. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-35, 51-57, and 90-97, of SEQ ID NO:6, respectively.

The DNA sequence of the variable region of the heavy chain of MAb 6-2 has been determined, and is presented in SEQ ID NO:7; the amino acid sequence encoded thereby is presented in SEQ ID NO:8. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 31-35, 50-66, and 99-107 of SEQ ID NO:8, respectively.

#### EXAMPLE 7: Assays for Measuring Loss of Barrier Function

A method provided herein involves use of IL-4 antagonists to inhibit IL-4-induced damage to epithelium, including but not limited to lung epithelium or intestinal epithelium. Damage to epithelium can result in loss of barrier function. A number of techniques are known for determining whether an epithelial layer is intact. The following are examples of techniques that may be employed in assessing the ability of an IL-4 antagonist to inhibit IL-4-induced damage to epithelium and loss of epithelial barrier function.

Cells that may be employed in preparing *in vitro* models of epithelium (epithelial barriers) are known. For example, Calu-3 human lung epithelial cells are suitable for use in barrier function studies. Another suitable cell line is the human intestinal epithelial cell line designated T84. T84 cells are cultured under conditions that result in formation of a monolayer of epithelial cells on a permeable support, as described in Madara, J. and K. Dharmasathaphorn (*J. Cell Biol.*, 101:2124-2133, 1985), Madara, J. and J. Stafford (*J. Clin. Invest.* 83:724-727, 1989), and Youakim, A. and M. Ahdieh (*Am. J. Physiol.* 276 (*Gastrointest. Liver Physiol.* 39):G1279-G1288, 1999). The cultured monolayers are tested for properties such as resistance to passive transepithelial ion flow (such resistance indicating an intact monolayer performing a barrier function). The thus-generated epithelial monolayer simulates the intestinal epithelial barrier.

One type of assay determines whether a particular radiolabeled compound is able to cross an epithelial monolayer (e.g., a monolayer generated as described above). Transport of the radiolabeled compound across the monolayer indicates that the barrier is permeable rather than intact. One such procedure is mannitol flux analysis, which assesses movement of radiolabeled mannitol (e.g., <sup>3</sup>H mannitol) across a monolayer (see Madara and Stafford, *supra*).

Methods for imaging a monolayer are identified in Madara and Stafford, *supra*. Such imaging methods are an alternative for assessing the condition of an epithelial layer, after exposure to IL-4 with or without an antagonist.

Youakim and Ahdieh, *supra*, discuss proteins that are part of "tight junction" complexes in intact intestinal epithelial barriers, and report studies of the effect of IFN- $\gamma$  on proteins associated with tight junctions. Other techniques for studying the effect of a cytokine on barrier function are described. For example, the effect of a cytokine on monolayer permeability may be assessed by transepithelial electrical resistance measurements, using techniques described in the reference.



- U.S. Patent 6,033,688 also describes procedures that may be employed in studies of barrier permeability; see especially examples 1 and 4 of the patent. Human tracheal epithelial cells were cultured under conditions that yielded a monolayer exhibiting transepithelial electrical resistance. Transepithelial resistance (indicating an intact barrier) was determined using a voltmeter. The effect of a particular reagent (HGH) on the epithelial monolayer was assessed by exposing the monolayer to HGH, and then measuring ion transport activities in Ussing chambers, by standard methods (column 8, lines 40-56). Similar studies were conducted on monolayers that were generated from bronchial epithelial cells from a human cystic fibrosis patient (example 4, column 11).
- Using any of the above-described barrier function assay procedures, an epithelial monolayer is exposed to IL-4 alone, or exposed to IL-4 in the presence of an IL-4 antagonist. The antagonist's ability to inhibit the IL-4-induced reduction in barrier function thus is assessed.
- In one such assay, a monolayer of T84 cells served as an *in vitro* model of an intestinal epithelial barrier, as discussed above. IL-4 added to the basolateral side of polarized epithelial cells was found to reduce barrier function by 70% within 48-72 hours of treatment. When an IL-4 receptor polypeptide was added at the same time as IL-4, the reduction in barrier function was prevented, and the barrier was maintained at the same level as untreated (control) cells. A soluble human IL-4 receptor polypeptide, consisting of the extracellular domain, was employed in the assay.
- The assay procedure also was conducted on a monolayer derived from lung epithelial cells, which served as an *in vitro* model of a lung epithelial barrier. IL-4 added to the basolateral side of polarized lung epithelial cells was found to reduce barrier function by 50% within 48-72 hours of treatment. When the IL-4 receptor polypeptide was added at the same time as IL-4, the reduction in barrier function was prevented, and the barrier was maintained at the same level as untreated (control) cells.

#### EXAMPLE 8: Monoclonal Antibody designated 12B5

- A human monoclonal antibody directed against human IL-4 receptor was prepared by the following procedure. The monoclonal antibody, which is designated 12B5, is a blocking antibody that functions as an IL-4 antagonist and as an IL-13 antagonist.
- The procedure began with immunization of a transgenic mouse with a soluble human IL-4 receptor polypeptide. Mouse strain ((CMD)++; (JKD)++; (HCo7)11952+/+; (KCo5)9272+/+), described in example 3 above, was employed. The antigen for immunization was purified soluble IL-4R, comprising the extracellular domain of human IL-4 receptor (500ug/ml). The mouse was initially immunized with 50 ug antigen emulsified in Complete Freund's Adjuvant, followed by two more immunizations with Incomplete Freund's Adjuvant at 50ug and then 25ug. Immunization was every two weeks by intraperitoneal injection. A specific human IgG anti-IL-4R titer from the serum was performed by ELISA, eight days after the last injection. A good titer against the target was

detected, and the mouse then was IV/IP boosted 25ug each on day -3 (i.e., 3 days before the mouse was sacrificed) and 15ug IV on day -2.

The mouse was sacrificed, and spleen cells were extracted and fused with the murine myeloma cell line P3x63Ag8.653 (ATCC CRL 1580). A conventional PEG fusion protocol was followed. The fusion was screened for HulgG and HuKappa, followed by a rescreen for human gamma and kappa specific to IL-4R. The positive clones were evaluated, and clone 12B5 was identified as a blocking antibody. The clone was subcloned; a hybridoma cell line that produces MAb 12B5 was isolated; and MAb 12B5 was purified from the supernatant.

12B5 was determined to be an IgG1 antibody, and to be fully human. Antibodies of other subclasses, such as IgG4 or IgM monoclonal antibodies, may be derived from 12B5. Techniques for altering (switching) the subclass/isotype of an antibody are known. The constant region of 12B5 may be replaced, for example, with a constant region derived from a human IgG4 antibody. Sequence information for a human IgG4 heavy chain is presented, for example, in Ellison et al. (*DNA Vol. 1*, no. 1, pp 11-18, 1981), which is hereby incorporated by reference herein.

DNA encoding the variable region of the light chain of MAb 12B5 was isolated, and the nucleotide sequence thereof was determined. The DNA sequence for the light chain variable region is presented as SEQ ID NO:9; the amino acid sequence encoded thereby is presented in SEQ ID NO:10. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-35, 51-57, and 90-99, of SEQ ID NO:10, respectively.

DNA encoding the variable region of the heavy chain of MAb 12B5 was isolated, and the nucleotide sequence thereof was determined. The DNA sequence for the heavy chain variable region is presented as SEQ ID NO:11; the amino acid sequence encoded thereby is presented in SEQ ID NO:12. CDR-1 of the heavy chain is believed to correspond to amino acids 31-35; CDR-2 to amino acids 50-65; and CDR-3 to amino acids 98-104 of SEQ ID NO:12.

#### **EXAMPLE 9: Additional Monoclonal Antibodies that inhibit both IL-4 and IL-13**

Additional human monoclonal antibodies were raised against human IL-4 receptor, by immunizing transgenic mice with a soluble human IL-4R polypeptide. The transgenic mice employed were selected from the transgenic mouse strains described in example 3.

Hybridoma cell lines secreting human monoclonal antibodies that specifically bind human IL-4R, and which are capable of functioning as IL-4 antagonists and IL-13 antagonists, were identified and isolated. The MAbs are designated 27A1, 5A1, and 63. Another MAb, designated 1B7, was derived from MAb 63, and differs from the parent antibody only in the light chain. 1B7 retains the ability to bind IL-4R and to function as an IL-4 antagonist and an IL-13 antagonist.

The DNA sequence of the variable region of the light chain of MAb 27A1 is presented in SEQ ID NO:13, and the encoded amino acid sequence is presented in SEQ ID NO:14. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-35, 51-57, and 90-99, of SEQ ID NO:14, respectively.

The DNA sequence for the variable region of the heavy chain of MAb 27A1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 31-35, 50-66, and 99-105, of SEQ ID NO:16, respectively.

- 5       The DNA sequence of the variable region of the light chain of MAb 5A1 is presented in SEQ ID NO:17, and the encoded amino acid sequence is presented in SEQ ID NO:18. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-34, 50-56, and 89-97 of SEQ ID NO:18, respectively.

- 10       The DNA sequence for the variable region of the heavy chain of MAb 5A1 is presented as SEQ ID NO:19, and the encoded amino acid sequence is presented in SEQ ID NO:20. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 31-35, 50-65, and 98-112 of SEQ ID NO:20, respectively.

- 15       The DNA sequence of the variable region of the light chain of MAb 63 is presented in SEQ ID NO:21, and the encoded amino acid sequence is presented in SEQ ID NO:22. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-34, 50-56, and 89-97 of SEQ ID NO:22, respectively.

- 20       The DNA sequence for the variable region of the heavy chain of MAb 63 is presented as SEQ ID NO:23, and the encoded amino acid sequence is presented in SEQ ID NO:24. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 31-35, 50-66, and 99-106 of SEQ ID NO:24, respectively.

- 25       The DNA sequence of the variable region of the light chain of MAb 1B7 is presented in SEQ ID NO:25, and the encoded amino acid sequence is presented in SEQ ID NO:26. Complementarity determining regions 1 to 3 (CDR 1-3) are believed to correspond to amino acids 24-34, 50-56, and 89-97 of SEQ ID NO:26, respectively.

- MAb 1B7 was derived from MAb 63, and the heavy chains of the two MAbs are identical. Thus, the DNA sequence for the variable region of the heavy chain of MAb 1B7 is presented as SEQ ID NO:23, and the encoded amino acid sequence is presented in SEQ ID NO:24. Complementarity determining regions 1 to 3 (CDRs 1-3) are believed to correspond to amino acids 31-35, 50-66, and 99-106 of SEQ ID NO:24, respectively.

**CLAIMS**

What is claimed is:

1. A human antibody that binds human IL-4 receptor (IL-4R), wherein said antibody is capable of inhibiting an IL-4-induced biological activity.
2. An antibody of claim 1, wherein said antibody additionally is capable of inhibiting an IL-13-induced biological activity.
3. An antibody of claim 2, wherein the light chain of said antibody comprises an amino acid sequence selected from the amino acid sequence presented in SEQ ID NO:6, SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:22, or SEQ ID NO:26.
4. An antibody of claim 2, wherein the heavy chain of said antibody comprises an amino acid sequence selected from the amino acid sequence presented in SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:16, SEQ ID NO:20, or SEQ ID NO:24.
5. A method of inhibiting an IL-4-induced biological activity *in vivo*, comprising administering an antibody of claim 1 to a human.
6. A method of inhibiting an IL-4-induced biological activity and an IL-13-induced biological activity *in vivo*, comprising administering an antibody of claim 2 to a human.
7. A method of claim 5, wherein said antibody is administered in the form of a composition that additionally comprises a diluent, excipient, or carrier.
8. A method of claim 6, wherein said antibody is administered in the form of a composition that additionally comprises a diluent, excipient, or carrier.
9. A method for treating septic arthritis, comprising administering an IL-4 antagonist to a mammal afflicted with septic arthritis.
10. A method for treating septic arthritis, comprising administering an antibody of claim 1 to a human afflicted with septic arthritis.
11. A method of claim 9, wherein said antagonist is a soluble form of the human IL-4 receptor (IL-4R) of SEQ ID NO:2, and the mammal is a human.

12. A method of claim 9, wherein said antagonist is an antibody that inhibits the binding of IL-4 to an IL-4R.
13. A method of claim 9, wherein said antagonist is administered in the form of a composition that additionally comprises a diluent, excipient, or carrier.

**FIGURE 1A**

ATG	GGG	TGG	CTT	TGC	TCT	GGG	CTC	CTG	TTC	CCT	GTG	AGC	TGC	CTG	-31
Met	Gly	Trp	Leu	Cys	Ser	Gly	Leu	Leu	Phe	Pro	Val	Ser	Cys	Leu	-11
GTC	CTG	CTG	CAG	GTG	GCA	AGC	TCT	GGG	AAC	ATG	AAG	GTC	TTG	CAG	15
Val	Leu	Leu	Gln	Val	Ala	Ser	Ser	Gly	Asn	Met	Lys	Val	Leu	Gln	5
GAG	CCC	ACC	TGC	GTC	TCC	GAC	TAC	ATG	AGC	ATC	TCT	ACT	TGC	GAG	60
Glu	Pro	Thr	Cys	Val	Ser	Asp	Tyr	Met	Ser	Ile	Ser	Thr	Cys	Glu	20
TGG	AAG	ATG	AAT	GGT	CCC	ACC	AAT	TGC	AGC	ACC	GAG	CTC	CGC	CTG	105
Trp	Lys	Met	Asn	Gly	Pro	Thr	Asn	Cys	Ser	Thr	Glu	Leu	Arg	Leu	35
TTG	TAC	CAG	CTG	GTT	TTT	CTG	CTC	TCC	GAA	GCC	CAC	ACG	TGT	ATC	150
Leu	Tyr	Gln	Leu	Val	Phe	Leu	Leu	Ser	Glu	Ala	His	Thr	Cys	Ile	50
CCT	GAG	AAC	AAC	GGA	GGC	GCG	GGG	TGC	GTG	TGC	CAC	CTG	CTC	ATG	195
Pro	Glu	Asn	Asn	Gly	Gly	Ala	Gly	Cys	Val	Cys	His	Leu	Leu	Met	65
GAT	GAC	GTG	GTC	AGT	GCG	GAT	AAC	TAT	ACA	CTG	GAC	CTG	TGG	GCT	240
Asp	Asp	Val	Val	Ser	Ala	Asp	Asn	Tyr	Thr	Leu	Asp	Leu	Trp	Ala	80
GGG	CAG	CAG	CTG	CTG	TGG	AAG	GGC	TCC	TTC	AAG	CCC	AGC	GAG	CAT	285
Gly	Gln	Gln	Leu	Leu	Trp	Lys	Gly	Ser	Phe	Lys	Pro	Ser	Glu	His	95
GTG	AAA	CCC	AGG	GCC	CCA	GGA	AAC	CTG	ACA	GTT	CAC	ACC	AAT	GTC	330
Val	Lys	Pro	Arg	Ala	Pro	Gly	Asn	Leu	Thr	Val	His	Thr	Asn	Val	110
TCC	GAC	ACT	CTG	CTG	CTG	ACC	TGG	AGC	AAC	CCG	TAT	CCC	CCT	GAC	375
Ser	Asp	Thr	Leu	Leu	Leu	Thr	Trp	Ser	Asn	Pro	Tyr	Pro	Pro	Asp	125
AAT	TAC	CTG	TAT	AAT	CAT	CTC	ACC	TAT	GCA	GTC	AAC	ATT	TGG	AGT	420
Asn	Tyr	Leu	Tyr	Asn	His	Leu	Thr	Tyr	Ala	Val	Asn	Ile	Trp	Ser	140
GAA	AAC	GAC	CCG	GCA	GAT	TTC	AGA	ATC	TAT	AAC	GTG	ACC	TAC	CTA	465
Glu	Asn	Asp	Pro	Ala	Asp	Phe	Arg	Ile	Tyr	Asn	Val	Thr	Tyr	Leu	155
GAA	CCC	TCC	CTC	CGC	ATC	GCA	GCC	AGC	ACC	CTG	AAG	TCT	GGG	ATT	510
Glu	Pro	Ser	Leu	Ser	Ile	Ala	Ala	Ser	Thr	Leu	Lys	Ser	Gly	Ile	170
TCC	TAC	AGG	GCA	CGG	GTG	AGG	GCC	TGG	GCT	CAG	TGC	TAT	AAC	ACC	555
Ser	Tyr	Arg	Ala	Arg	Val	Arg	Ala	Trp	Ala	Gln	Cys	Tyr	Asn	Thr	185
ACC	TGG	AGT	GAG	TGG	AGC	CCC	AGC	ACC	AAG	TGG	CAC	AAC	TCC	TAC	600
Thr	Trp	Ser	Glu	Trp	Ser	Pro	Ser	Thr	Lys	Trp	His	Asn	Ser	Tyr	200
AGG	GAG	CCC	TTC	GAG	CAG	CAC	CTC	CTG	CTG	GGC	GTC	AGC	GTT	TCC	645
Arg	Glu	Pro	Phe	Glu	Gln	His	Leu	Leu	Leu	Gly	Val	Ser	Val	Ser	215
TGC	ATT	GTC	ATC	CTG	GCC	GTC	TGC	CTG	TTG	TGC	TAT	GTC	AGC	ATC	690
Cys	Ile	Val	Ile	Leu	Ala	Val	Cys	Leu	Leu	Cys	Tyr	Val	Ser	Ile	230
ACC	AAG	ATT	AAG	AAA	GAA	TGG	TGG	GAT	CAG	ATT	CCC	AAC	CCA	GCC	735
Thr	Lys	Ile	Lys	Lys	Glu	Trp	Trp	Asp	Gln	Ile	Pro	Asn	Pro	Ala	245

**FIGURE 1B**

CGC	AGC	CGC	CTC	GTG	GCT	ATA	ATA	ATC	CAG	GAT	GCT	CAG	GGG	TCA	780
Arg	Ser	Arg	Leu	Val	Ala	Ile	Ile	Ile	Gln	Asp	Ala	Gln	Gly	Ser	260
CAG	TGG	GAG	AAG	CGG	TCC	CGA	GGC	CAG	GAA	CCA	GCC	AAG	TGC	CCA	825
Gln	Trp	Glu	Lys	Arg	Ser	Arg	Gly	Gln	Glu	Pro	Ala	Lys	Cys	Pro	275
CAC	TGG	AAG	AAT	TGT	CTT	ACC	AAG	CTC	TTG	CCC	TGT	TTT	CTG	GAG	870
His	Trp	Lys	Asn	Cys	Leu	Thr	Lys	Leu	Leu	Pro	Cys	Phe	Leu	Glu	290
CAC	AAC	ATG	AAA	AGG	GAT	GAA	GAT	CCT	CAC	AAG	GCT	GCC	AAA	GAG	915
His	Asn	Met	Lys	Arg	Asp	Glu	Asp	Pro	His	Lys	Ala	Ala	Lys	Glu	305
ATG	CCT	TTC	CAG	GGC	TCT	GGA	AAA	TCA	GCA	TGG	TGC	CCA	GTG	GAG	960
Met	Pro	Phe	Gln	Gly	Ser	Gly	Lys	Ser	Ala	Trp	Cys	Pro	Val	Glu	320
ATC	AGC	AAG	ACA	GTC	CTC	TGG	CCA	GAG	AGC	ATC	AGC	GTG	GTG	CGA	1005
Ile	Ser	Lys	Thr	Val	Leu	Trp	Pro	Glu	Ser	Ile	Ser	Val	Val	Arg	335
TGT	GTG	GAG	TTG	TTT	GAG	GCC	CCG	GTG	GAG	TGT	GAG	GAG	GAG	GAG	1050
Cys	Val	Glu	Leu	Phe	Glu	Ala	Pro	Val	Glu	Cys	Glu	Glu	Glu	Glu	350
GAG	GTA	GAG	GAA	GAA	AAA	GGG	AGC	TTC	TGT	GCA	TCG	CCT	GAG	AGC	1095
Glu	Val	Glu	Glu	Glu	Lys	Gly	Ser	Phe	Cys	Ala	Ser	Pro	Glu	Ser	365
AGC	AGG	GAT	GAC	TTC	CAG	GAG	GGA	AGG	GAG	GGC	ATT	GTG	GCC	CGG	1140
Ser	Arg	Asp	Asp	Phe	Gln	Glu	Gly	Arg	Glu	Gly	Ile	Val	Ala	Arg	380
CTA	ACA	GAG	AGC	CTG	TTC	CTG	GAC	CTG	CTC	GGA	GAG	GAG	AAT	GGG	1185
Leu	Thr	Glu	Ser	Leu	Phe	Leu	Asp	Leu	Leu	Gly	Glu	Glu	Asn	Gly	395
GGC	TTT	TGC	CAG	CAG	GAC	ATG	GGG	GAG	TCA	TGC	CTT	CTT	CCA	CCT	1230
Gly	Phe	Cys	Gln	Gln	Asp	Met	Gly	Glu	Ser	Cys	Leu	Leu	Pro	Pro	410
TCG	GGA	AGT	ACG	AGT	GCT	CAC	ATG	CCC	TGG	GAT	GAG	TTC	CCA	AGT	1275
Ser	Gly	Ser	Thr	Ser	Ala	His	Met	Pro	Trp	Asp	Glu	Phe	Pro	Ser	425
GCA	GGG	CCC	AAG	GAG	GCA	CCT	CCC	TGG	GGC	AAG	GAG	CAG	CCT	CTC	1320
Ala	Gly	Pro	Lys	Glu	Ala	Pro	Pro	Trp	Gly	Lys	Glu	Gln	Pro	Leu	440
CAC	CTG	GAG	CCA	AGT	CCT	CCT	GCC	AGC	CCG	ACC	CAG	AGT	CCA	GAC	1365
His	Leu	Glu	Pro	Ser	Pro	Pro	Ala	Ser	Pro	Thr	Gln	Ser	Pro	Asp	455
AAC	CTG	ACT	TGC	ACA	GAG	ACG	CCC	CTC	GTC	ATC	GCA	GGC	AAC	CCT	1410
Asn	Leu	Thr	Cys	Thr	Glu	Thr	Pro	Leu	Val	Ile	Ala	Gly	Asn	Pro	470
GCT	TAC	CGC	AGC	TTC	AGC	AAC	TCC	CTG	AGC	CAG	TCA	CCG	TGT	CCC	1455
Ala	Tyr	Arg	Ser	Phe	Ser	Asn	Ser	Leu	Ser	Gln	Ser	Pro	Cys	Pro	485
AGA	GAG	CTG	GGT	CCA	GAC	CCA	CTG	CTG	GCC	AGA	CAC	CTG	GAG	GAA	1500
Arg	Glu	Leu	Gly	Pro	Asp	Pro	Leu	Leu	Ala	Arg	His	Leu	Glu	Glu	500
GTA	GAA	CCC	GAG	ATG	CCC	TGT	GTC	CCC	CAG	CTC	TCT	GAG	CCA	ACC	1545
Val	Glu	Pro	Glu	Met	Pro	Cys	Val	Pro	Gln	Leu	Ser	Glu	Pro	Thr	515

**FIGURE 1C**

ACT	GTG	CCC	CAA	CCT	GAG	CCA	GAA	ACC	TGG	GAG	CAG	ATC	CTC	CGC	1590
Thr	Val	Pro	Gln	Pro	Glu	Pro	Glu	Thr	Trp	Glu	Gln	Ile	Leu	Arg	530
CGA	AAT	GTC	CTC	CAG	CAT	GGG	GCA	GCT	GCA	GCC	CCC	GTC	TCG	GCC	1635
Arg	Asn	Val	Leu	Gln	His	Gly	Ala	Ala	Ala	Ala	Pro	Val	Ser	Ala	545
CCC	ACC	AGT	GGC	TAT	CAG	GAG	TTT	GTA	CAT	GCG	GTG	GAG	CAG	GGT	1680
Pro	Thr	Ser	Gly	Tyr	Gln	Glu	Phe	Val	His	Ala	Val	Glu	Gln	Gly	560
GGC	ACC	CAG	GCC	AGT	GCG	GTG	GTG	GGC	TTG	GGT	CCC	CCA	GGA	GAG	1725
Gly	Thr	Gln	Ala	Ser	Ala	Val	Val	Gly	Leu	Gly	Pro	Pro	Gly	Glu	575
GCT	GGT	TAC	AAG	GCC	TTC	TCA	AGC	CTG	CTT	GCC	AGC	AGT	GCT	GTG	1770
Ala	Gly	Tyr	Lys	Ala	Phe	Ser	Ser	Leu	Leu	Ala	Ser	Ser	Ala	Val	590
TCC	CCA	GAG	AAA	TGT	GGG	TTT	GGG	GCT	AGC	AGT	GGG	GAA	GAG	GGG	1815
Ser	Pro	Glu	Lys	Cys	Gly	Phe	Gly	Ala	Ser	Ser	Gly	Glu	Glu	Gly	605
TAT	AAG	CCT	TTC	CAA	GAC	CTC	ATT	CCT	GGC	TGC	CCT	GGG	GAC	CCT	1860
Tyr	Lys	Pro	Phe	Gln	Asp	Leu	Ile	Pro	Gly	Cys	Pro	Gly	Asp	Pro	620
GCC	CCA	GTC	CCT	GTC	CCC	TTG	TTC	ACC	TTT	GGA	CTG	GAC	AGG	GAG	1905
Ala	Pro	Val	Pro	Val	Pro	Leu	Phe	Thr	Phe	Gly	Leu	Asp	Arg	Glu	635
CCA	CCT	CGC	AGT	CCG	CAG	AGC	TCA	CAT	CTC	CCA	AGC	AGC	TCC	CCA	1950
Pro	Pro	Arg	Ser	Pro	Gln	Ser	Ser	His	Leu	Pro	Ser	Ser	Ser	Pro	650
GAG	CAC	CTG	GGT	CTG	GAG	CCG	GGG	GAA	AAG	GTA	GAG	GAC	ATG	CCA	1995
Glu	His	Leu	Gly	Leu	Glu	Pro	Gly	Glu	Lys	Val	Glu	Asp	Met	Pro	665
AAG	CCC	CCA	CTT	CCC	CAG	GAG	CAG	GCC	ACA	GAC	CCC	CTT	GTG	GAC	2040
Lys	Pro	Pro	Leu	Pro	Gln	Glu	Gln	Ala	Thr	Asp	Pro	Leu	Val	Asp	680
AGC	CTG	GGC	AGT	GGC	ATT	GTC	TAC	TCA	GCC	CTT	ACC	TGC	CAC	CTG	2085
Ser	Leu	Gly	Ser	Gly	Ile	Val	Tyr	Ser	Ala	Leu	Thr	Cys	His	Leu	695
TGC	GGC	CAC	CTG	AAA	CAG	TGT	CAT	GGC	CAG	GAG	GAT	GGT	GGC	CAG	2130
Cys	Gly	His	Leu	Lys	Gln	Cys	His	Gly	Gln	Glu	Asp	Gly	Gly	Gln	710
ACC	CCT	GTC	ATG	GCC	AGT	CCT	TGC	TGT	GGC	TGC	TGC	TGT	GGA	GAC	2175
Thr	Pro	Val	Met	Ala	Ser	Pro	Cys	Cys	Gly	Cys	Cys	Cys	Gly	Asp	725
AGG	TCC	TCG	CCC	CCT	ACA	ACC	CCC	CTG	AGG	GCC	CCA	GAC	CCC	TCT	2220
Arg	Ser	Ser	Pro	Pro	Thr	Thr	Pro	Leu	Arg	Ala	Pro	Asp	Pro	Ser	740
CCA	GGT	GGG	GTT	CCA	CTG	GAG	GCC	AGT	CTG	TGT	CCG	GCC	TCC	CTG	2265
Pro	Gly	Gly	Val	Pro	Leu	Glu	Ala	Ser	Leu	Cys	Pro	Ala	Ser	Leu	755
GCA	CCC	TCG	GGC	ATC	TCA	GAG	AAG	AGT	AAA	TCC	TCA	TCA	TCC	TTC	2310
Ala	Pro	Ser	Gly	Ile	Ser	Glu	Lys	Ser	Lys	Ser	Ser	Ser	Ser	Phe	770
CAT	CCT	GCC	CCT	GGC	AAT	GCT	CAG	AGC	TCA	AGC	CAG	ACC	CCC	AAA	2355
His	Pro	Ala	Pro	Gly	Asn	Ala	Gln	Ser	Ser	Ser	Gln	Thr	Pro	Lys	785
ATC	GTG	AAC	TTT	GTC	TCC	GTG	GGA	CCC	ACA	TAC	ATG	AGG	GTC	TCT	2400
Ile	Val	Asn	Phe	Val	Ser	Val	Gly	Pro	Thr	Tyr	Met	Arg	Val	Ser	800







FIGURE 2C

## FIGURE 3A

AATTAGCGGC	CGCTGTCGAC	AAGCTTCGAA	TTCAGTATCG	ATGTGGGGTA	50
CCTACTGTCC	CGGGATTGCG	GATCCGCGAT	GATATCGTTG	ATCCTCGAGT	100
GCGGCCGCGAC	TATGCAAAAA	AAAGCCCGCT	CATTAGGCGG	GCTCTTGGCA	150
GAACATATCC	ATCGCGTCCG	CCATCTCCAG	CAGCCGCGACG	CGCGCGCATCT	200
CGGGCAGCGT	TGGGTCTCTG	CCACGGGTGC	GCATGATCGT	GCTCCTGTCTG	250
TTGAGGACCC	GGCTAGGCTG	GCGGGGTTCG	CTTACTGGTT	AGCAGAAATGA	300
ATCACCGATA	CGCGAGCGAA	CGTGAAGCGA	CTGCTGCTGC	AAAACGTCGT	350
CGACCTGAGC	AACAACATGA	ATGGTCTTCG	GTTTCCGTGT	TTCGTAAAGT	400
CTGGAACACG	GGAAGTCAGC	GCCCTGCACC	ATTATGTTCC	GGATCTGCAT	450
CGCAGGATGC	TGCTGGCTAC	CCTGTGGAAC	ACCTACATCT	GTATTAAACGA	500
AGCGCTGGCA	TTGACCCCTGA	GTGATTTTTC	TCTGGTCCCG	CCGCATCCAT	550
ACCGCCAGTT	GTTTACCCTC	ACAACGTTCC	AGTAACCGGG	CATGTTTCATC	600
ATCAGTAACC	CGTATCGTGA	GCATCCTCTC	TCGTTTTCATC	GGTATCATTAA	650
CCCCCATGAA	CAGAAATTC	CCCTTACACG	GAGGCATCAA	GTGACCAAAC	700
AGGAAAAAAC	CGCCCTTAAC	ATGGCCCGCT	TTATCAGAAG	CCAGACATTA	750
ACGCTTCTTG	AGAAACTCAA	CGAGCTGGAC	GCGGATGAAC	AGGCAGACAT	800
CTTGTGAATCG	GTCGAGACC	ACGCTGATGA	GCTTTACCGC	AGCTGCCCTCG	850
CGCGTTTCGG	TGATGACGGT	GAAAACCTCT	GACACATGCA	GCTCCCGGAG	900
ACGGTCACAG	CTTGTCTGTA	AGCGGATGCC	GGGAGCAGAC	AAGCCCGTCA	950
GGGCGCGTCA	GCGGGTGTTC	GCGGGTGTTC	GGGCGCAGCC	ATGACCCAGT	1000
CACGTAGCGA	TAGCGGAGTG	TATACTGGCT	TAACTATGCG	GCATCAGAG	1050
AGATTGTACT	GAGAGTGCAC	CATATGCGGT	GTGAAATACC	GCACAGATGC	1100
GTAAGGAGAA	AATACCGCAT	CAGGCGCTCT	TCCGCTTCCCT	CGCTCACTGA	1150
CTCGCTGCGC	TCGGTTCGTT	GGCTGCGGCG	AGCGGTATCA	GCTCACTCAA	1200
AGGCGGTAAT	ACGGTTATCC	ACAGAATCAG	GGGATAACGC	AGGAAAGAAC	1250
ATGTGAGCAA	AAGGCCAGCA	AAAGGCCAGG	AACCGTAAAA	AGGCCGCGTT	1300
GCTGGCGTTT	TTCCATAGGC	TCCGCCCCCC	TGACGAGCAT	CACAAAAATC	1350
GACGCTCAAG	TCAGAGGTGG	CGAAACCCGA	CAGGACTATA	AAGATACCAG	1400
GCGTTTCCCC	CTGGAAGCTC	CCTCGTGCGC	TCTCCTGTTC	CGACCCCTGCC	1450
GCTTACCNGA	TACCTGTCCG	CCTTTCCTCC	TTCCGGGAAG	GTGGCGGCTTT	1500
CTCATAGCTC	ACGCTGTAGG	TATCTCAGTT	CGGTGTAGGT	CGTTCGCTCC	1550
AAGCTGGGCT	GTGTGCACGA	ACCCCCCGTT	CAGCCCCGAC	GCTCGGCGCTT	1600
ATCCGGTAAC	TATCGTCTTG	AGTCCAACCC	GTAAGACAC	GACTTATFCG	1650
CACCTGGCAGC	AGCCAGGCGC	GCCTTGGCCT	AAGAGGCCAC	TGGTAAACAGG	1700
ATTAGCAGAG	CGAGGTATGT	AGGCGGTGCT	ACAGAGTTCT	TGAAGTGGTG	1750
GCCTAACTAC	GCTCTACACTA	GAAGGACAGT	ATTGGTATC	TGCGCTCTGC	1800
TGAAGCCAGT	TACCTTCGGA	AAAAGAGTTG	GTAGCTCTTG	ATCCGGGCAA	1850
CAAACACCAG	CTGGTAGCGG	TGGTTTTTTT	GTTTGCAAGC	AGCAGATTAC	1900
GCGCAGAAAA	AAAGGATCTC	AAGAAGATCC	TTTGATCTTT	TCTACGGGGT	1950
CTGACGCTCA	TGGGAACGAA	AACTCACGTT	AAGGGATTTT	GGTCAATGAA	2000
TTATCAAAAA	GGATCTTCAC	CTAGATCCTT	TTAAATTTAA	AATGAAGTTT	2050
TAAATCAATC	TAAAGTATAT	ATGAGTAAAC	TTGGTCTGAC	AGTTACCAAT	2100
GCTTAATCAG	TGAGGCCACCT	ATCTCAGCGA	TCTGTCTATT	TCGTTTATCC	2150
ATAGTTGCCT	GACTTCCCGT	CGTGTAGATA	ACTACGATAC	GGGAGGGGCTT	2200
ACCATCTGGC	CCAGTGCTG	CAATGATACC	GCGAGACCCA	CGCTCACCGG	2250
CTCCAGATTT	ATCAGCAATA	AACCAGCCAG	CCGGAAGGGC	CGAGCGCGCA	2300
AGTGGTCCCT	CAACTTTATC	CGCCTCCATC	CAGTCTATTA	ATTGTTGCCG	2350
GGAGCTAGA	GTAAGTAGTT	CGCCAGTTAA	TAGTTTGCGC	AACGTTGTTG	2400

## FIGURE 3B

CCATTGCTGC	AGGCATCGTG	GTGTCACGCT	CGTCGTTGG	TATGGCTTCA	2450
TTCAGCTCCG	GTTCCCAACG	ATCAAGGCGA	GTTACATGAT	CCCCCATGTT	2500
GTGCAAAAAA	GCGGTTAGCT	CCTTCGGTCC	TCCGATCGTT	GTCAGAAAGTA	2550
AGTTGGCCGC	AGTGTTATCA	CTCATGGTTA	TGGCAGCACT	GCATAATTCT	2600
CTTACTGTCA	TGCCATCCGT	AAGATGCTTT	TCTGTGACTG	GTGAGTACTC	2650
AACCAAGTCA	TTCTGAGAAT	AGTGTATGCG	GCGACCGAGT	TGCTCTTGCC	2700
CGGCGTCAAC	ACGGGATAAT	ACCGCGCCAC	ATAGCAGAAC	TTTAAAAGTG	2750
CTCATCATTG	GAAAACGTTT	TTCGGGGCGA	AAACTCTCAA	GGATCTTACC	2800
GCTGTTGAGA	TCCAGTTCGA	TGTAACCCAC	TCGTGCACCC	AACTGATCTT	2850
CAGCATCTTT	TACTTTCACC	AGCGTTTCTG	GGTGAGCAAA	AACAGGAAGG	2900
CAAAATGCCG	CAAAAAAGGG	AATAAGGGCG	ACACGGAAT	GTTGAATACT	2950
CATACTCTTC	CTTTTTCAAT	ATTATTGAAG	CATTATTCAG	GGTTATTGTC	3000
TCATGAGCGG	ATACATATTT	GAATGTATTT	AGAAAAATAA	ACAAATAGGG	3050
GTTCGCGCA	CATTTCCCCG	AAAAGTGCCA	CCTGACGTCT	AAGAAACCAT	3100
TATTATCATG	ACATTAACTT	ATAAAAATAG	GCGTATCACG	AGGCCCTTTC	3150
GTCTTCAAG					3159

## FIGURE 4

1B7Vk	EIVLTQSPSS	VSASVGRVT	ITCRASQGIS	N.RLAWYOOK	PGKAPKLLIY	IASILQRGVP
63Vk	D-QM-----	-----	-----	T-W-----H-	-----	V-S--S--
12B5Vk	-----GT	L-L-P-E-A-	LS-----SV-	SSY-----	--Q-R--F	G--SRAT-I-
6-2Vk	-----GT	L-L-P-E-A-	LS-----SV-	TSY-----	--Q-R--	G--SRAT-I-
5A1Vk	-----AT	L-L-P-E-A-	LS-----SV-	.SY-----	--Q-R--	H--NRAT-I-
27A1Vk	-----GT	L-L-P-E-A-	LS-----SV-	SSY-----	--Q-R--	G--SRAT-I-
1B7Vk	SRSFGSGGT	DFITITRLQ	PEDEATYYCQ	Q.ANSFPFTF	GGTKVDIK	
63Vk	-----	-----SS--	-----	-----	-----	
12B5Vk	D-----	-----S-E	-----V-----	-YGS-P-W--	-Q-----	E--
6-2Vk	D-----	-----S-E	-----V-----	-YGS-P--	-Q-----	E--
5A1Vk	A-----	-----SS-E	-----V-----	-RSNW-L--	-G-----	E--
27A1Vk	D-----	-----S-E	-----V-----	-YGS-P-W--	-Q-----	E--

FIGURE 5

1B7VH	EYQVLESGN	LIVQEGGSLRL	SCAASGTFS	SVAMSWRQA	PCKGLEWVSS	ITGGSGSTYY	ADSVKGRFTI	SRDNS
#63VH	---	---	---	---	---	---	---	---
12BSVH	--LVQ--	G-H-	-G-	RN-R-	-G-	-T-A-N-	-A	-A
6-2VH	Q-IV---	G V-	-IQ-	-A-	-AV	-SYD-SKK-	-A	-A
5A1VH	--LVQ--	G-H-	T-G-	NFV-H-	-T-	-G-	-A	-A
27A1VH	Q-LV---	G V-R-	-S-	R-G-H-	-R-	-WFE-NNQ-	-V	-V
1B7VH	KNTLYLQMS	LRAEDTAIFY	CAKDN.R.G.	. . . FFHYWG	QGLTLTVSS			
#63VH	---	---	---	---	---			
12BSVH	--S----	M-VY-	-RGR.-	--YV-D-	---			
6-2VH	---	VY-	-REG-R-	--S-D-	---			
5A1VH	--SS----	M-VY-	R-REMRG	VII DY-D-	---			
27A1VH	---E---	VY-	-R.GKY.	--Y-D-	---			

## SEQUENCE LISTING

&lt;110&gt; Pluenneke, John

&lt;120&gt; USE OF INTERLEUKIN-4 ANTAGONISTS AND COMPOSITIONS THEREOF

&lt;130&gt; 3005-WO

&lt;140&gt; --to be assigned--

&lt;141&gt; 2001-05-25

&lt;150&gt; 09/579,808

&lt;151&gt; 2000-05-26

&lt;150&gt; 09/665,343

&lt;151&gt; 2000-09-19

&lt;150&gt; 09/785,934

&lt;151&gt; 2001-02-15

&lt;150&gt; 09/847,816

&lt;151&gt; 2001-05-01

&lt;160&gt; 26

&lt;170&gt; PatentIn version 3.0

&lt;210&gt; 1

&lt;211&gt; 2478

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(2475)

&lt;220&gt;

&lt;221&gt; mat\_peptide

&lt;222&gt; (76)..()

&lt;400&gt; 1

atg	ggg	tgg	ctt	tgc	tct	ggg	ctc	ctg	ttc	cct	gtg	agc	tgc	ctg	gtc	48
Met	Gly	Trp	Leu	Cys	Ser	Gly	Leu	Leu	Phe	Pro	Val	Ser	Cys	Leu	Val	
-25					-20					-15					-10	

ctg	ctg	cag	gtg	gca	agc	tct	ggg	aac	atg	aag	gtc	ttg	cag	gag	ccc	96
Leu	Leu	Gln	Val	Ala	Ser	Ser	Gly	Asn	Met	Lys	Val	Leu	Gln	Glu	Pro	
			-5					-1	1				5			

acc	tgc	gtc	tcc	gac	tac	atg	agc	atc	tct	act	tgc	gag	tgg	aag	atg	144
Thr	Cys	Val	Ser	Asp	Tyr	Met	Ser	Ile	Ser	Thr	Cys	Glu	Trp	Lys	Met	
		10					15					20				

aat	ggt	ccc	acc	aat	tgc	agc	acc	gag	ctc	cgc	ctg	ttg	tac	cag	ctg	192
Asn	Gly	Pro	Thr	Asn	Cys	Ser	Thr	Glu	Leu	Arg	Leu	Leu	Tyr	Gln	Leu	
	25					30					35					

gtt	ttt	ctg	ctc	tcc	gaa	gcc	cac	acg	tgt	atc	cct	gag	aac	aac	gga	240
Val	Phe	Leu	Leu	Ser	Glu	Ala	His	Thr	Cys	Ile	Pro	Glu	Asn	Asn	Gly	
40				45					50						55	

ggc gcg ggg tgc	gtg tgc cac ctg	ctc atg gat gac	gtg gtc agt gcg	288
Gly Ala Gly Cys	Val Cys His Leu	Leu Met Asp Asp	Val Val Ser Ala	
	60	65	70	
gat aac tat aca	ctg gac ctg tgg	gct ggg cag cag	ctg ctg tgg aag	336
Asp Asn Tyr Thr	Leu Asp Leu Trp	Ala Gly Gln Gln	Leu Leu Trp Lys	
	75	80	85	
ggc tcc ttc aag	ccc agc gag cat	gtg aaa ccc agg	gcc cca gga aac	384
Gly Ser Phe Lys	Pro Ser Glu His	Val Lys Pro Arg	Ala Pro Gly Asn	
	90	95	100	
ctg aca gtt cac	acc aat gtc tcc	gac act ctg ctg	ctg acc tgg agc	432
Leu Thr Val His	Thr Asn Val Ser	Asp Thr Leu Leu	Leu Thr Trp Ser	
	105	110	115	
aac ccg tat ccc	cct gac aat tac	ctg tat aat cat	ctc acc tat gca	480
Asn Pro Tyr Pro	Pro Asp Asn Tyr	Leu Tyr Asn His	Leu Thr Tyr Ala	
	120	125	130	135
gtc aac att tgg	agt gaa aac gac	ccg gca gat ttc	aga atc tat aac	528
Val Asn Ile Trp	Ser Glu Asn Asp	Pro Ala Asp Phe	Arg Ile Tyr Asn	
	140	145	150	
gtg acc tac cta	gaa ccc tcc ctc	cgc atc gca gcc	agc acc ctg aag	576
Val Thr Tyr Leu	Glu Pro Ser Leu	Arg Ile Ala Ala	Ser Thr Leu Lys	
	155	160	165	
tct ggg att tcc	tac agg gca cgg	gtg agg gcc tgg	gct cag tgc tat	624
Ser Gly Ile Ser	Tyr Arg Ala Arg	Val Arg Ala Trp	Ala Gln Cys Tyr	
	170	175	180	
aac acc acc tgg	agt gag tgg agc	ccc agc acc aag	tgg cac aac tcc	672
Asn Thr Thr Trp	Ser Glu Trp Ser	Pro Ser Thr Lys	Trp His Asn Ser	
	185	190	195	
tac agg gag ccc	ttc gag cag cac	ctc ctg ctg ggc	gtc agc gtt tcc	720
Tyr Arg Glu Pro	Phe Glu Gln His	Leu Leu Leu Gly	Val Ser Val Ser	
	200	205	210	215
tgc att gtc atc	ctg gcc gtc tgc	ctg ttg tgc tat	gtc agc atc acc	768
Cys Ile Val Ile	Leu Ala Val Cys	Leu Leu Cys Tyr	Val Ser Ile Thr	
	220	225	230	
aag att aag aaa	gaa tgg tgg gat	cag att ccc aac	cca gcc cgc agc	816
Lys Ile Lys Lys	Glu Trp Trp Asp	Gln Ile Pro Asn	Pro Ala Arg Ser	
	235	240	245	
cgc ctc gtg gct	ata ata atc cag	gat gct cag ggg	tca cag tgg gag	864
Arg Leu Val Ala	Ile Ile Ile Gln	Asp Ala Gln Gly	Ser Gln Trp Glu	
	250	255	260	
aag cgg tcc cga	ggc cag gaa cca	gcc aag tgc cca	cac tgg aag aat	912
Lys Arg Ser Arg	Gly Gln Glu Pro	Ala Lys Cys Pro	His Trp Lys Asn	
	265	270	275	
tgt ctt acc aag	ctc ttg ccc tgt	ttt ctg gag cac	aac atg aaa agg	960
Cys Leu Thr Lys	Leu Leu Pro Cys	Phe Leu Leu His	Asn Met Lys Arg	
	280	285	290	295

gat gaa gat cct cac aag gct gcc aaa gag atg cct ttc cag ggc tct Asp Glu Asp Pro His Lys Ala Ala Lys Glu Met Pro Phe Gln Gly Ser 300 305 310	1008
gga aaa tca gca tgg tgc cca gtg gag atc agc aag aca gtc ctc tgg Gly Lys Ser Ala Trp Cys Pro Val Glu Ile Ser Lys Thr Val Leu Trp 315 320 325	1056
cca gag agc atc agc gtg gtg cga tgt gtg gag ttg ttt gag gcc ccg Pro Glu Ser Ile Ser Val Val Arg Cys Val Glu Leu Phe Glu Ala Pro 330 335 340	1104
gtg gag tgt gag gag gag gag gta gag gaa gaa aaa ggg agc ttc Val Glu Cys Glu Glu Glu Glu Val Glu Glu Lys Gly Ser Phe 345 350 355	1152
tgt gca tcg cct gag agc agc agg gat gac ttc cag gag gga agg gag Cys Ala Ser Pro Glu Ser Ser Arg Asp Asp Phe Gln Glu Gly Arg Glu 360 365 370 375	1200
ggc att gtg gcc cgg cta aca gag agc ctg ttc ctg gac ctg ctc gga Gly Ile Val Ala Arg Leu Thr Glu Ser Leu Phe Leu Asp Leu Leu Gly 380 385 390	1248
gag gag aat ggg ggc ttt tgc cag cag gac atg ggg gag tca tgc ctt Glu Glu Asn Gly Gly Phe Cys Gln Gln Asp Met Gly Glu Ser Cys Leu 395 400 405	1296
ctt cca cct tcg gga agt acg agt gct cac atg ccc tgg gat gag ttc Leu Pro Pro Ser Gly Ser Thr Ser Ala His Met Pro Trp Asp Glu Phe 410 415 420	1344
cca agt gca ggg ccc aag gag gca cct ccc tgg ggc aag gag cag cct Pro Ser Ala Gly Pro Lys Glu Ala Pro Pro Trp Gly Lys Glu Gln Pro 425 430 435	1392
ctc cac ctg gag cca agt cct cct gcc agc ccg acc cag agt cca gac Leu His Leu Glu Pro Ser Pro Pro Ala Ser Pro Thr Gln Ser Pro Asp 440 445 450 455	1440
aac ctg act tgc aca gag acg ccc ctc gtc atc gca ggc aac cct gct Asn Leu Thr Cys Thr Glu Thr Pro Leu Val Ile Ala Gly Asn Pro Ala 460 465 470	1488
tac cgc agc ttc agc aac tcc ctg agc cag tca ccg tgt ccc aga gag Tyr Arg Ser Phe Ser Asn Ser Leu Ser Gln Ser Pro Cys Pro Arg Glu 475 480 485	1536
ctg ggt cca gac cca ctg ctg gcc aga cac ctg gag gaa gta gaa ccc Leu Gly Pro Asp Pro Leu Leu Ala Arg His Leu Glu Glu Val Glu Pro 490 495 500	1584
gag atg ccc tgt gtc ccc cag ctc tct gag cca acc act gtg ccc caa Glu Met Pro Cys Val Pro Gln Leu Ser Glu Pro Thr Thr Val Pro Gln 505 510 515	1632
cct gag cca gaa acc tgg gag cag atc ctc cgc cga aat gtc ctc cag Pro Glu Pro Glu Thr Trp Glu Gln Ile Leu Arg Arg Asn Val Leu Gln 520 525 530 535	1680



cat ggg gca gct gca gcc ccc gtc tgc gcc ccc acc agt ggc tat cag His Gly Ala Ala His Ala Pro Val Ser Ala Gly Thr Ser Gly Tyr Gln 540 545 550	1728
gag ttt gta cat gcg gtg gag cag ggt ggc acc cag gcc agt gcg gtg Glu Phe Val His Ala Val Glu Gln Gly Gly Thr Gln Ala Ser Ala Val 555 560 565	1776
gtg ggc ttg ggt ccc cca gga gag gct ggt tac aag gcc ttc tca agc Val Gly Leu Gly Pro Pro Gly Glu Ala Gly Tyr Lys Ala Phe Ser Ser 570 575 580	1824
ctg ctt gcc agc agt gct gtg tcc cca gag aaa tgt ggg ttt ggg gct Leu Leu Ala Ser Ser Ala Val Ser Pro Glu Lys Cys Gly Phe Gly Ala 585 590 595	1872
agc agt ggg gaa gag ggg tat aag cct ttc caa gac ctc att cct ggc Ser Ser Gly Glu Glu Gly Tyr Lys Pro Phe Gln Asp Leu Ile Pro Gly 600 605 610 615	1920
tgc cct ggg gac cct gcc cca gtc cct gtc ccc ttg ttc acc ttt gga Cys Pro Gly Asp Pro Ala Pro Val Pro Val Pro Leu Phe Thr Phe Gly 620 625 630	1968
ctg gac agg gag cca cct cgc agt ccg cag agc tca cat ctc cca agc Leu Asp Arg Glu Pro Pro Arg Ser Pro Gln Ser Ser His Leu Pro Ser 635 640 645	2016
agc tcc cca gag cac ctg ggt ctg gag ccg ggg gaa aag gta gag gac Ser Ser Pro Glu His Leu Gly Leu Glu Pro Gly Glu Lys Val Glu Asp 650 655 660	2064
atg cca aag ccc cca ctt ccc cag gag cag gcc aca gac ccc ctt gtg Met Pro Lys Pro Pro Leu Pro Gln Glu Gln Ala Thr Asp Pro Leu Val 665 670 675	2112
gac agc ctg ggc agt ggc att gtc tac tca gcc ctt acc tgc cac ctg Asp Ser Leu Gly Ser Gly Ile Val Tyr Ser Ala Leu Thr Cys His Leu 680 685 690 695	2160
tgc ggc cac ctg aaa cag tgt cat ggc cag gag gat ggt ggc cag acc Cys Gly His Leu Lys Gln Cys His Gly Gln Glu Asp Gly Gly Gln Thr 700 705 710	2208
cct gtc atg gcc agt cct tgc tgt ggc tgc tgc tgt gga gac agg tcc Pro Val Met Ala Ser Pro Cys Cys Gly Cys Cys Gly Asp Arg Ser 715 720 725	2256
tgc ccc cct aca acc ccc ctg agg gcc cca gac ccc tct cca ggt ggg Ser Pro Pro Thr Thr Pro Leu Arg Ala Pro Asp Pro Ser Pro Gly Gly 730 735 740	2304
gtt cca ctg gag gcc agt ctg tgt ccg gcc tcc ctg gca ccc tgc ggc Val Pro Leu Glu Ala Ser Leu Cys Pro Ala Ser Leu Ala Pro Ser Gly 745 750 755	2352
atc tca gag aag agt aaa tcc tca tca tcc ttc cat cct gcc cct ggc Ile Ser Glu Lys Ser Lys Ser Ser Ser Ser Phe His Pro Ala Pro Gly 760 765 770 775	2400

aat gct cag agc tca agc cag acc ccc aaa atc gtg aac ttt gtc tcc 2448  
 Asn Ala Gln Ser Ser Ser Gln Thr Pro Lys Ile Val Asn Phe Val Ser  
                     780                    785                    790

gtg gga ccc aca tac atg agg gtc tct tat 2478  
 Val Gly Pro Thr Tyr Met Arg Val Ser  
                     795                    800

<210> 2  
 <211> 825  
 <212> PRT  
 <213> Homo sapiens

<400> 2

Met Gly Trp Leu Cys Ser Gly Leu Leu Phe Pro Val Ser Cys Leu Val  
 -25                    -20                    -15                    -10

Leu Leu Gln Val Ala Ser Ser Gly Asn Met Lys Val Leu Gln Glu Pro  
                     -5                    -1 1                    5

Thr Cys Val Ser Asp Tyr Met Ser Ile Ser Thr Cys Glu Trp Lys Met  
                     10                    15                    20

Asn Gly Pro Thr Asn Cys Ser Thr Glu Leu Arg Leu Leu Tyr Gln Leu  
                     25                    30                    35

Val Phe Leu Leu Ser Glu Ala His Thr Cys Ile Pro Glu Asn Asn Gly  
                     40                    45                    50                    55

Gly Ala Gly Cys Val Cys His Leu Leu Met Asp Asp Val Val Ser Ala  
                     60                    65                    70

Asp Asn Tyr Thr Leu Asp Leu Trp Ala Gly Gln Gln Leu Leu Trp Lys  
                     75                    80                    85

Gly Ser Phe Lys Pro Ser Glu His Val Lys Pro Arg Ala Pro Gly Asn  
                     90                    95                    100

Leu Thr Val His Thr Asn Val Ser Asp Thr Leu Leu Leu Thr Trp Ser  
                     105                    110                    115

Asn Pro Tyr Pro Pro Asp Asn Tyr Leu Tyr Asn His Leu Thr Tyr Ala  
                     120                    125                    130                    135

Val Asn Ile Trp Ser Glu Asn Asp Pro Ala Asp Phe Arg Ile Tyr Asn  
                     140                    145                    150

Val Thr Tyr Leu Glu Pro Ser Leu Arg Ile Ala Ala Ser Thr Leu Lys  
 155 160 165

Ser Gly Ile Ser Tyr Arg Ala Arg Val Arg Ala Trp Ala Gln Cys Tyr  
 170 175 180

Asn Thr Thr Trp Ser Glu Trp Ser Pro Ser Thr Lys Trp His Asn Ser  
 185 190 195

Tyr Arg Glu Pro Phe Glu Gln His Leu Leu Leu Gly Val Ser Val Ser  
 200 205 210 215

Cys Ile Val Ile Leu Ala Val Cys Leu Leu Cys Tyr Val Ser Ile Thr  
 220 225 230

Lys Ile Lys Lys Glu Trp Trp Asp Gln Ile Pro Asn Pro Ala Arg Ser  
 235 240 245

Arg Leu Val Ala Ile Ile Ile Gln Asp Ala Gln Gly Ser Gln Trp Glu  
 250 255 260

Lys Arg Ser Arg Gly Gln Glu Pro Ala Lys Cys Pro His Trp Lys Asn  
 265 270 275

Cys Leu Thr Lys Leu Leu Pro Cys Phe Leu Glu His Asn Met Lys Arg  
 280 285 290 295

Asp Glu Asp Pro His Lys Ala Ala Lys Glu Met Pro Phe Gln Gly Ser  
 300 305 310

Gly Lys Ser Ala Trp Cys Pro Val Glu Ile Ser Lys Thr Val Leu Trp  
 315 320 325

Pro Glu Ser Ile Ser Val Val Arg Cys Val Glu Leu Phe Glu Ala Pro  
 330 335 340

Val Glu Cys Glu Glu Glu Glu Glu Val Glu Glu Glu Lys Gly Ser Phe  
 345 350 355

Cys Ala Ser Pro Glu Ser Ser Arg Asp Asp Phe Gln Glu Gly Arg Glu  
 360 365 370 375

Gly Ile Val Ala Arg Leu Thr Glu Ser Leu Phe Leu Asp Leu Leu Gly  
 380 385 390

Glu Glu Asn Gly Gly Phe Cys Gln Gln Asp Met Gly Glu Ser Cys Leu  
 395 400 405

Leu Pro Pro Ser Gly Ser Thr Ser Ala His Met Pro Trp Asp Glu Phe  
 410 415 420

Pro Ser Ala Gly Pro Lys Glu Ala Pro Pro Trp Gly Lys Glu Gln Pro  
 425 430 435

Leu His Leu Glu Pro Ser Pro Pro Ala Ser Pro Thr Gln Ser Pro Asp  
 440 445 450 455

Asn Leu Thr Cys Thr Glu Thr Pro Leu Val Ile Ala Gly Asn Pro Ala  
 460 465 470

Tyr Arg Ser Phe Ser Asn Ser Leu Ser Gln Ser Pro Cys Pro Arg Glu  
 475 480 485

Leu Gly Pro Asp Pro Leu Leu Ala Arg His Leu Glu Glu Val Glu Pro  
 490 495 500

Glu Met Pro Cys Val Pro Gln Leu Ser Glu Pro Thr Thr Val Pro Gln  
 505 510 515

Pro Glu Pro Glu Thr Trp Glu Gln Ile Leu Arg Arg Asn Val Leu Gln  
 520 525 530 535

His Gly Ala Ala Ala Ala Pro Val Ser Ala Pro Thr Ser Gly Tyr Gln  
 540 545 550

Glu Phe Val His Ala Val Glu Gln Gly Gly Thr Gln Ala Ser Ala Val  
 555 560 565

Val Gly Leu Gly Pro Pro Gly Glu Ala Gly Tyr Lys Ala Phe Ser Ser  
 570 575 580

Leu Leu Ala Ser Ser Ala Val Ser Pro Glu Lys Cys Gly Phe Gly Ala  
 585 590 595

Ser Ser Gly Glu Glu Gly Tyr Lys Pro Phe Gln Asp Leu Ile Pro Gly  
 600 605 610 615

Cys Pro Gly Asp Pro Ala Pro Val Pro Val Pro Leu Phe Thr Phe Gly  
 620 625 630

Leu Asp Arg Glu Pro Pro Arg Ser Pro Gln Ser Ser His Leu Pro Ser  
 635 640 645  
 Ser Ser Pro Glu His Leu Gly Leu Glu Pro Gly Glu Lys Val Glu Asp  
 650 655 660  
 Met Pro Lys Pro Pro Leu Pro Gln Glu Gln Ala Thr Asp Pro Leu Val  
 665 670 675  
 Asp Ser Leu Gly Ser Gly Ile Val Tyr Ser Ala Leu Thr Cys His Leu  
 680 685 690 695  
 Cys Gly His Leu Lys Gln Cys His Gly Gln Glu Asp Gly Gly Gln Thr  
 700 705 710  
 Pro Val Met Ala Ser Pro Cys Cys Gly Cys Cys Cys Gly Asp Arg Ser  
 715 720 725  
 Ser Pro Pro Thr Thr Pro Leu Arg Ala Pro Asp Pro Ser Pro Gly Gly  
 730 735 740  
 Val Pro Leu Glu Ala Ser Leu Cys Pro Ala Ser Leu Ala Pro Ser Gly  
 745 750 755  
 Ile Ser Glu Lys Ser Lys Ser Ser Ser Ser Phe His Pro Ala Pro Gly  
 760 765 770 775  
 Asn Ala Gln Ser Ser Ser Gln Thr Pro Lys Ile Val Asn Phe Val Ser  
 780 785 790  
 Val Gly Pro Thr Tyr Met Arg Val Ser  
 795 800

<210> 3  
 <211> 8  
 <212> PRT  
 <213> Artificial

<220>  
 <223> FLAG peptide

<400> 3

Asp Tyr Lys Asp Asp Asp Asp Lys  
 1 5

<210> 4  
 <211> 3159  
 <212> DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; "Cloning vector pGPik"

&lt;400&gt; 4

aattagcggc cgctgtcgac aagcttcgaa ttcagtatcg atgtggggta cctactgtcc	60
cgggattcgc gatccgcgat gatatcgtt atcctcagat gcggccgcag tatgcaaaaa	120
aaagcccgct cattaggcgg gctcttggca gaacatatcc atcgcgtccg ccatctccag	180
cagccgcacg cggcgcatct cgggcagcgt tgggtccttg ccacgggtgc gcgatgcgt	240
gctcctgtcg ttgaggaccg ggctaggctg gcgggggttg cttactgtgt agcagaatga	300
atcaccgata cgcgagcgaa cgtgaagcga ctgctgctgc aaaacgtctg cgacctgagc	360
aacaacatga atggctcttc gtttccgtgt ttcgtaaaagt ctggaaacgc ggaagtcagc	420
gccctgcacc attatgttcc ggatctgcgt cgcaggatgc tgctggctac cctgtggaac	480
acctacatct gtattaacga agcgcctggca ttgacctga gtgatttttc tctgtgcccg	540
ccgcatecat acgcgcagtt gtttaccctc acaacgttcc agtaaccggg catgttctac	600
atcagtaacc cgtatcgtga gcatacctctc tcgtttctac ggatcatta cccccatgaa	660
cagaaattcc cccttacacg gaggcaccaa gtgaccaaac aggaaaaaac cgccttaac	720
atggcccgct tttatcagaag ccagacatta acgcttctgg agaaactcaa cgagctggac	780
gcggatgaac aggcagacat ctgtgaatcg cttcacgacc acgctgatga gctttaccgc	840
agctgcctcg cgcgtttcgg tgatgacggc gaaaacctct gacacatgca gctcccgag	900
acggctcacg cttgtctgta agcggatgcc gggagcagac aagcccgta cggcgcgta	960
gcgggtgtgt gcgggtgtcg gggcgacgcc atgaccagtc cactgtagca tagcggagtg	1020
tatactggct taactatcgc gcatacagac agattgtact gagagtgcac catatgcggt	1080
gtgaaatacc gcacagatgc gtaaggagaa aataccgcgt caggcgctct tccgttctc	1140
cgtcactga ctcgctgcgc tcggctgttc ggctgcggcg agcgggtatca gctcactcaa	1200
aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac atgtgagcaa	1260
aaggccagca aaaggccagg aaccgtaaaa aggcgcgcgt gctggcgctt ttcctaggg	1320
tccgcccccc tgaccgacat cacaaaaatc gacgctcaag tcagaggttg cgaaccgga	1380
caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc	1440
cgaccctgcc gcttaccgga tacctgtccg cttttctccc ttcgggaagc gtggcgcttt	1500
ctcatagctc acgctgtagg tatctcagtt cggtgtaggc cgttcgctcc aagctgggct	1560
gtgtgcacga accccccgtt cagcccgacc gctgcgcctt atccggtaac tatcgtcttg	1620
agtccaaccc ggtaagacac gacttatcgc cactggcagc agccaggcgc gccttggcct	1680

```

aagagggcac  tggtaacagg  attagcagag  cgaggatgt  aggggtgtct  acagagttct  1740
tgaagtgggt  gctaacctac  ggctacacta  gaaggacagt  atttgggtatc  tgcgtctctgc  1800
tgaagccagt  taccttcgga  aaaagagttg  gtgctcttg  atccggcaaaa  caaaccaccg  1860
ctggtagcgg  tggttttttt  gtttgcaagc  agcagattac  gcgcagaaaa  aaaggatctc  1920
aagaagatcc  tttgatcttt  tctacggggt  ctgacgtctc  gtggaacgaa  aactcacgtt  1980
aagggatttt  ggcatgaga  ttatcaaaaa  ggatcttcac  ctagatcctt  ttaaatataa  2040
aatgaagttt  taaatcaatc  taaagtatat  atgagtaaac  ttggtctgac  agttaccaat  2100
gcttaatcag  tgaggcacct  atctcagcga  tctgtctatt  tcgttcaccc  atagttgcct  2160
gactcccggt  cgtgtagata  actacgatac  gggagggcct  accatctggc  ccagtgctg  2220
caatgatacc  gcgagacca  cgctcacgg  ctccagattt  atcagcaata  aaccagccag  2280
ccggaagggg  cgagcgaga  agtggctctg  caactttatc  cgctccatc  cagtctatta  2340
attgttgccg  ggaagctaga  gtaagtagtt  cgccagtaa  tagtttgcgc  aacgttggtg  2400
ccattgctgc  aggcacgtg  gtgtcacgct  cgtcgtttgg  tatggcttca  ttcagctcgc  2460
gttccaacg  atcaaggcga  gttacatgat  ccccatgtt  gtgcaaaaaa  gcgggttagct  2520
ccttcggctc  tccgatcgtt  gtcagaagta  agttggccgc  agtgttatca  ctcatgggta  2580
tggcagcact  gcataattct  ctactgtca  tgccatccgt  aagatgcttt  tctgtgactg  2640
gtgagtactc  aaccaagtc  ttctgagaat  agtgatgctg  gcgaccgagt  tgcctctgcc  2700
cggcgctaac  acgggataat  accgcgccac  atagcagaac  tttaaaagt  ctcatcattg  2760
gaaaaagttc  ttcggggcga  aaactctcaa  ggatcttacc  gctgttgaga  tccagttcga  2820
tgtaaccacc  tcgtgcacc  aactgatctt  cagcatcttt  tactttcacc  agcgtttctg  2880
ggtagcaaaa  aacaggaagg  caaatgccg  caaaaaagg  aataaggcgc  acacggaaat  2940
gttgaatact  catactcttc  ctttttcaat  attattgaag  catttatcag  ggttattgtc  3000
tcattgagcgg  atacatattt  gaatgtattt  agaaaaataa  acaaataggg  gttccgcgca  3060
catttccccc  aaaagtcca  cctgacgtct  aagaaacct  tattatcatg  acattaacct  3120
ataaaaatag  gcgtatcacg  aggccttttc  gtcttcaag  3159

```

```

<210> 5
<211> 321
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (1)..(321)

```

<400> 5  
gaa att gtg ttg acg cag tct cca ggc acc ctg tct ttg tct cca ggg 48  
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
1 5 10 15

gaa aga gcc acc ctc tcc tgc agg gcc agt cag agt gtt agc acc agc 96  
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Thr Ser  
20 25 30

tac tta gcc tgg tac cag cag aaa cct ggc cag gct ccc agg ctc ctc 144  
Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
35 40 45

atc tat ggt gca tcc agc agg gcc act ggc atc cca gac agg ttc agt 192  
Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
50 55 60

ggc agt ggg tct ggg aca gac ttc act ctc acc atc agc aga ctg gag 240  
Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
65 70 75 80

cct gaa gat ttt gca gtg tat tac tgt cag cag tat ggt agt tca ccg 288  
Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
85 90 95

acg ttc ggc caa ggg acc aag gtg gaa atc aaa 321  
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
100 105

<210> 6  
<211> 107  
<212> PRT  
<213> Homo sapiens

<400> 6  
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Thr Ser  
20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
85 90 95



Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
 100 105

<210> 7  
 <211> 354  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(354)

<400> 7  
 cag gtg cag ctg gtg gag tct ggg gga ggc gtg gtc cag cct ggg agg 48  
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15  
 tcc ctg aga ctc tcc tgt gca gcc tct gga ttc gcc ttc agt agc tat 96  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ser Tyr  
 20 25 30  
 gct att cag tgg gtc cgc cag gct cca ggc aag ggg ctg gag tgg gtg 144  
 Ala Ile Gln Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45  
 gca gtt ata tca tat gat gga agc aag aaa tac tat gca gac tcc gtg 192  
 Ala Val Ile Ser Tyr Asp Gly Ser Lys Lys Tyr Tyr Ala Asp Ser Val  
 50 55 60  
 aag ggc cga ttc acc atc tcc aga gac aat tcc aag aat acg ctg tat 240  
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80  
 ttg caa atg aac agc ctg aga gct gag gac acg gct gtg tat tac tgt 288  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 gcg aga gag ggg aga cgt ggg tcg ttt gac tac tgg ggc cag gga acc 336  
 Ala Arg Glu Gly Arg Arg Gly Ser Phe Asp Tyr Trp Gly Gln Gly Thr  
 100 105 110  
 ctg gtc acc gtc tcc tca 354  
 Leu Val Thr Val Ser Ser  
 115

<210> 8  
 <211> 118  
 <212> PRT  
 <213> Homo sapiens

<400> 8  
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ser Tyr  
 20 25 30

Ala Ile Gln Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Lys Lys Tyr Tyr Ala Asp Ser Val  
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg Glu Gly Arg Arg Gly Ser Phe Asp Tyr Trp Gly Gln Gly Thr  
100 105 110

Leu Val Thr Val Ser Ser  
115

<210> 9  
<211> 327  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (1)..(327)

<400> 9  
gaa att gtg ttg acg cag tct cca ggc acc ctg tct ttg tct cca ggg 48  
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
1 5 10 15  
gaa aga gcc acc ctc ttc tgc agg gcc agt cag agt gtt agc agc agc 96  
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser  
20 25 30  
tac tta gcc tgg tac cag cag aaa cct gcc cag gct ccc agg ctc ctc 144  
Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
35 40 45  
atc ttt ggt gca tcc agc agg gcc act gcc atc cca gac agg ttc agt 192  
Ile Phe Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
50 55 60  
ggc agt ggg tct ggg aca gac ttc act ctc acc atc agc aga ctg gag 240  
Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
65 70 75 80  
cct gaa gat ttt gca gtg tat tac tgt cag cag tat ggt agc tca cct 288  
Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
85 90 95

ccg tgg acg ttc ggc caa ggg acc aag gtg gaa atc aaa 327  
 Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
                   100                  105  
  
 <210> 10  
  
 <211> 109  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 10  
  
 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
 1                  5                  10                  15  
  
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser  
                   20                  25                  30  
  
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
                   35                  40                  45  
  
 Ile Phe Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
                   50                  55                  60  
  
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
 65                  70                  75                  80  
  
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
                   85                  90                  95  
  
 Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
                   100                  105  
  
  
 <210> 11  
 <211> 345  
 <212> DNA  
 <213> Homo sapiens  
  
 <220>  
 <221> CDS  
 <222> (1)..(345)  
  
 <400> 11  
 gag gtt cag ctg gtg cag tct ggg gga ggc ttg gta cat cct ggg ggg 48  
 Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1                  5                  10                  15  
  
 tcc ctg aga ctc tcc tgt gca ggc tct gga ttc acc ttc agt aga aat 96  
 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Arg Asn  
                   20                  25                  30  
  
 gct atg ttc tgg gtt cgc cag gct cca gga aaa ggt ctg gag tgg gta 144  
 Ala Met Phe Thr Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                  40                  45

tca ggt att ggt act ggt ggt gcc aca aac tat gca gac tcc gtg aag 192  
 Ser Gly Ile Gly Thr Gly Gly Ala Thr Asn Tyr Ala Asp Ser Val Lys  
 50 55 60

ggc cga ttc acc atc tcc aga gac aat gcc aag aac tcc ttg tat ctt 240  
 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu  
 65 70 75 80

caa atg aac agc ctg aga gcc gag gac atg gct gtg tat tac tgt gca 288  
 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

aga ggg agg tac tac ttt gac tac tgg ggc cag gga acc ctg gtc acc 336  
 Arg Gly Arg Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr  
 100 105 110

gtc tcc tca 345  
 Val Ser Ser  
 115

<210> 12  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens

<400> 12

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Arg Asn  
 20 25 30

Ala Met Phe Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ser Gly Ile Gly Thr Gly Gly Ala Thr Asn Tyr Ala Asp Ser Val Lys  
 50 55 60

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu  
 65 70 75 80

Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

Arg Gly Arg Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr  
 100 105 110

Val Ser Ser  
 115

<210> 13  
 <211> 327  
 <212> DNA  
 <213> Homo sapiens

<220>

<221> CDS  
 <222> (1)..(327)

<400> 13  
 gaa att gtg ttg acg cag tct cca ggc acc ctg tct ttg tct cca ggg 48  
 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15  
 gaa aga gcc acc ctc tcc tgc agg gcc agt cag agt gtt agc agc agc 96  
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser  
 20 25 30  
 tac tta gcc tgg tac caa cag aaa cct ggc cag gct ccc agg ctc ctc 144  
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
 35 40 45  
 atc tat ggt gca tcc agc agg gcc act ggc atc cca gac agg ttc agt 192  
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
 50 55 60  
 ggc agt ggg tct ggg aca gac ttc act ctc acc atc agc aga ctg gag 240  
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
 65 70 75 80  
 cct gaa gat ttt gca gtg tat tac tgt cag cag tat ggt agc tca cct 288  
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
 85 90 95  
 ccg tgg acg ttc ggc caa ggg acc aag gtg gaa atc aaa 327  
 Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
 100 105

<210> 14  
 <211> 109  
 <212> PRT  
 <213> Homo sapiens

<400> 14

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15  
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser  
 20 25 30  
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu  
 35 40 45  
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser  
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro  
85 90 95

Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
100 105

<210> 15  
<211> 348  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (1)..(348)

<400> 15  
cag gtg cag ctg gtg gag tct ggg gga ggc gtg gtc cag cct ggg agg 48  
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
1 5 10 15

tcc ctg aga ctc tcc tgt tca gcg tct gga ttc acc ttc agt aga tat 96  
Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Arg Tyr  
20 25 30

ggc atg cac tgg gtc cgc cag gct cca ggc agg ggg ctg gag tgg gtg 144  
Gly Met His Trp Val Arg Gln Ala Pro Gly Arg Gly Leu Glu Trp Val  
35 40 45

gca att ata tgg ttt gaa gga aat aat caa tac tat gca gac tcc gtg 192  
Ala Ile Ile Trp Phe Glu Gly Asn Asn Gln Tyr Tyr Ala Asp Ser Val  
50 55 60

aag ggc cga ttc acc gtc tcc aga gac aat tcc aag aac acg ctg tat 240  
Lys Gly Arg Phe Thr Val Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80

ctg gaa atg aac agc ctg aga gcc gag gac acg gct gtg tat tac tgt 288  
Leu Glu Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95

gcg aga ggg aag tac tac ttt gac tac tgg ggc cag gga acc ctg gtc 336  
Ala Arg Gly Lys Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val  
100 105 110

acc gtc tcc tca 348  
Thr Val Ser Ser  
115

<210> 16  
<211> 116  
<212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ser Ala Ser Gly Phe Thr Phe Ser Arg Tyr  
 20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Arg Gly Leu Glu Trp Val  
 35 40 45

Ala Ile Ile Trp Phe Glu Gly Asn Asn Gln Tyr Tyr Ala Asp Ser Val  
 50 55 60

Lys Gly Arg Phe Thr Val Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

Leu Glu Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Ala Arg Gly Lys Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val  
 100 105 110

Thr Val Ser Ser  
 115

&lt;210&gt; 17

&lt;211&gt; 321

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(321)

&lt;400&gt; 17

gaa att gtg ttg aca cag tct cca gcc acc ctg tct ttg tct cca ggg 48  
 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15

gaa aga gcc acc ctc tcc tgc agg gcc agt cag agt gtt agc agc tac 96  
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr  
 20 25 30

tta gcc tgg tac caa cag aaa cct gcc cag gct ccc agg ctc ctc atc 144  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile  
 35 40 45

tat cat gca tcc aac agg gcc act gcc atc cca gcc agg ttc agt gcc 192  
 Tyr His Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly  
 50 55 60

agt ggg tct ggg aca gac ttc act ctc acc atc agc agc cta gag cct 240  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro  
 65 70 75 80

gaa gat ttt gca gtt tat tac tgt cag cag cgt agc aac tgg cct ctc 288  
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Leu  
 85 90 95

act ttc ggc gga ggg acc aag gtg gag atc aaa 321  
 Thr Phe Gly Gly Thr Lys Val Glu Ile Lys  
 100 105

<210> 18  
 <211> 107  
 <212> PRT  
 <213> Homo sapiens

<400> 18

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile  
 35 40 45

Tyr His Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly  
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro  
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Leu  
 85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
 100 105

<210> 19  
 <211> 369  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(369)

<400> 19



gag gtg caa ctg gtg cag tct ggg gga ggc ttg gta cat cct ggg ggg 48  
 Glu Val Gln Leu Val Gln Ser Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15

tcc ctg aga ctc acc tgt gca ggc tct gga ttc acc ttc agt aac ttt 96  
 Ser Leu Arg Leu Thr Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Phe  
 20 25 30

gtt atg cac tgg gtt cgc cag act cca gga caa ggt ctg gag tgg gtt 144  
 Val Met His Trp Val Arg Gln Thr Pro Gly Gln Gly Leu Glu Trp Val  
 35 40 45

tca gct att ggt act ggt ggt ggc aca tac tat gca gac tcc gtg aag 192  
 Ser Ala Ile Gly Thr Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys  
 50 55 60

ggc cga ttc acc atc tcc aga gac aat gcc aag agc tcc tta tat ctt 240  
 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Ser Ser Leu Tyr Leu  
 65 70 75 80

caa atg aac agc ctg aga gcc gag gac atg gct gtg tat tac tgt gca 288  
 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95

aga gat cgg cct atg gtt cgg gga gtc att ata gac tac ttt gac tac 336  
 Arg Asp Arg Pro Met Val Arg Gly Val Ile Ile Asp Tyr Phe Asp Tyr  
 100 105 110

tgg ggc cag gga acc ctg gtc acc gtc tcc tca 369  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> 20  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens

<400> 20

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Thr Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Phe  
 20 25 30

Val Met His Trp Val Arg Gln Thr Pro Gly Gln Gly Leu Glu Trp Val  
 35 40 45

Ser Ala Ile Gly Thr Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys  
 50 55 60

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Ser Ser Leu Tyr Leu  
 65 70 75 80

Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
85 95

Arg Asp Arg Pro Met Val Arg Gly Val Ile Ile Asp Tyr Phe Asp Tyr  
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
115 120

<210> 21  
<211> 321  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (1)..(321)

<400> 21  
gac atc cag atg acc cag tct cca tct tcc gtg tct gca tct gta gga 48  
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly  
1 5 10 15

gac aga gtc acc atc act tgt cgg gcg agt cag ggt att agc acc tgg 96  
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Thr Trp  
20 25 30

tta gcc tgg tat cag cat aaa cca ggg aaa gcc cct aag ctc ctg atc 144  
Leu Ala Trp Tyr Gln His Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
35 40 45

tat gtt gca tcc agt ttg caa agt ggg gtc cca tca agg ttc agc ggc 192  
Tyr Val Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60

agt gga tct ggg aca gat ttc act ctc acc atc agc agc ctg cag cct 240  
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
65 70 75 80

gaa gat ttt gca act tac tat tgt caa cag gct aat agt ttc cca ttc 288  
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe  
85 90 95

act ttc ggc cct ggg acc aaa gtg gat atc aaa 321  
Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys  
100 105

<210> 22  
<211> 107  
<212> PRT  
<213> Homo sapiens

<400> 22

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly  
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Thr Trp  
 20 25 30

Leu Ala Trp Tyr Gln His Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
 35 40 45

Tyr Val Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro  
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe  
 85 90 95

Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys  
 100 105

<210> 23

<211> 351

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(351)

<400> 23

gag gtg cag gtg ttg gag tcg ggg gga aac ttg gta cag cct ggg ggg 48  
 Glu Val Gln Val Leu Glu Ser Gly Gly Asn Leu Val Gln Pro Gly Gly  
 1 5 10 15

tcc ctg aga ctc tcc tgt gca gcc tct gga ttc acc ttt agc agc tat 96  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
 20 25 30

gcc atg agc tgg gtc cgc cag gct cca ggg aag ggg ctg gag tgg gtc 144  
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

tct tct att act ggt agt ggg ggt agc aca tac tac gca gac tcc gtg 192  
 Ser Ser Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val  
 50 55 60

aag ggc cgg ttc acc atc tcc aga gac aat tcc aag aac acg ctg tat 240  
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80

ctg caa atg aac agc ctg aga gcc gag gac acg gcc att ttt tac tgt 288  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Ile Phe Tyr Cys  
 85 90 95

gcg aaa gat aac cgg gga ttc ttt cac tat tgg ggc cag gga acc ctg 336  
 Ala Lys Asp Asn Arg Gly Phe Phe His Tyr Trp Gly Gln Gly Thr Leu  
 100 105 110

gtc acc gtc tcc tca  
Val Thr Val Ser Ser  
115

351

<210> 24  
<211> 117  
<212> PRT  
<213> Homo sapiens

<400> 24

Glu Val Gln Val Leu Glu Ser Gly Gly Asn Leu Val Gln Pro Gly Gly  
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45

Ser Ser Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val  
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Ile Phe Tyr Cys  
85 90 95

Ala Lys Asp Asn Arg Gly Phe Phe His Tyr Trp Gly Gln Gly Thr Leu  
100 105 110

Val Thr Val Ser Ser  
115

<210> 25  
<211> 321  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (1)..(321)

<400> 25

gaa att gtg ttg aca cag tct cca tct tcc gtg tct gca tct gta gga  
Glu Ile Val Leu Thr Gln Ser Pro Ser Val Ser Ala Ser Val Gly  
1 5 10 15

48

gac aga gtc acc atc act tgt cgg gcg agt cag ggt att agc aat cgg 96  
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Arg  
 20 25 30

tta gcc tgg tat cag cag aaa cca gga aaa gcc cct aaa ctc ctc atc 144  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
 35 40 45

tat att gca tcc att ttg caa agg ggg gtc cca tca aga ttc agc ggc 192  
 Tyr Ile Ala Ser Ile Leu Gln Arg Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

agt gga tct ggg aca gat ttc act ctc acc atc acc agg ctg cag cct 240  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Thr Arg Leu Gln Pro  
 65 70 75 80

gaa gat ttt gca acc tac tat tgt caa cag gca aac agt ttc cca ttc 288  
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe  
 85 90 95

act ttc ggc cct ggg acc aaa gtg gat atc aaa 321  
 Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys  
 100 105

<210> 26  
 <211> 107  
 <212> PRT  
 <213> Homo sapiens

<400> 26

Glu Ile Val Leu Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly  
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Arg  
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile  
 35 40 45

Tyr Ile Ala Ser Ile Leu Gln Arg Gly Val Pro Ser Arg Phe Ser Gly  
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Thr Arg Leu Gln Pro  
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe  
 85 90 95

Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys  
 100 105

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
WO 01/92340 A3

- (51) International Patent Classification<sup>7</sup>: C07K 16/28, A61K 39/395, A61P 19/02, 31/04, A61K 38/17
- (72) Inventor; and  
(75) Inventor/Applicant (for US only): PLUENNEKE, John, D. [US/US]; 5816 Hickory Place, Paréville, MO 64152 (US).
- (21) International Application Number: PCT/US01/17094
- (22) International Filing Date: 25 May 2001 (25.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/579,808 26 May 2000 (26.05.2000) US  
09/665,343 19 September 2000 (19.09.2000) US  
09/785,934 15 February 2001 (15.02.2001) US  
09/847,816 1 May 2001 (01.05.2001) US
- (74) Agent: ANDERSON, Kathryn, A.; 51 University Street, Seattle, WA 98101-2936 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

*[Continued on next page]*

(54) Title: USE OF INTERLEUKIN-4 ANTAGONISTS AND COMPOSITIONS THEREOF

ATG GGG TGG CTT TGC TCT GGG CTC CTC TTC CCT GTG AGC TGC CTG -31  
Met Gly Trp Trp Cys Ser Gly Leu Leu Phe Pro Val Ser Cys Leu -11

GTG CTG CTG CAG GTG GCA AGC TCT GGG AAC ATG AAG GTC TTG CAG 15  
Val Leu Leu Gln Val Ala Ser Ser Gly Asn Met Lys Val Leu Gln 5

GAG CCC AGC TGC GTC TGC GAG TAC ATG AGC ATC TCT ACT TGC GAG 60  
Glu Pro Thr Cys Val Ser Asp Tyr Met Ser Ile Ser Thr Cys Glu 20

TGG AAG ATG AAT GGT CCC AAC AAT TGC AGC ACC GAG CTC CGC CTG 105  
Trp Lys Met Asn Gly Pro Thr Asn Cys Ser Thr Glu Leu Arg Leu 35

TTG TAC CAG CTG GTT TTT CTG TCC GAA GCC CAC AGC TGT ATC 150  
Leu Tyr Gln Leu Val Phe Leu Leu Ser Glu Ala His Thr Cys Ile 50

CCT GAG AAC AAC GGA GGC GCG GAG TGC GTG TGC CAG CTG CTC ATG 195  
Pro Glu Asn Asn Gly Gly Ala Gly Cys Val Cys His Leu Leu Met 65

GAT GAC GTG GTC AGT GCG GAT AAC TAT ACA CTG CAG CTG TGG GCT 240  
Asp Asp Val Val Ser Ala Asp Asn Tyr Thr Leu Asp Leu Trp Ala 80

GCG CAG CAG CTG CTG TGG AAG GGC TCC TTC AAG CCC AGC GAG CAT 285  
Gly Gln Gln Leu Leu Trp Lys Gly Ser Phe Lys Pro Ser Glu His 95

GTG AAA CCC AGG GCC CCA GAA AAC CTG ACA GTT CAC ACC AAT GTC 330  
Val Lys Pro Arg Ala Pro Gly Gln Leu Thr Val His Thr Asn Val 110

TCC GAC ATT CTG CTG CTG ACC TGG AGC AAC CCG TAT CCC CCG GAC 375  
Ser Asp Thr Leu Leu Thr Thr Tyr Ser Asn Pro Tyr Pro Pro Asp 125

AAT TAC CTG TAT AAT CAT CTC ACC TAT GCA GTC AAC ATT TGG AGT 420  
Asn Tyr Leu Tyr Asn His Leu Thr Tyr Ala Val Asn Ile Trp Ser 140

GAA AAC GAC CCG GCA GAT TTC AGA ATC TAT AAC GTG ACC TAC CTA 465  
Glu Asn Asp Pro Ala Asp Phe Arg Ile Tyr Asn Val Thr Tyr Leu 155

GAA CCC TCC CTC CAC ATC GCA GCG AGC ACC CTG AAG TCT GGG ATT 510  
Glu Pro Ser Leu Arg Ile Ala Ala Ser Thr Leu Lys Ser Gly Ile 170

TCC TAC AAG CCA CAG GTG AGG GCG TGG GGT GAG TGC TAT AAC ACC 555  
Ser Tyr Arg Ala Arg Val Arg Ala Trp Ala Gln Cys Tyr Asn Thr 185

ACC TGG AAT GAG TGG AGC CTC AGC ACC AAG TGG CAG AAC TCC TAC 600  
Thr Trp Ser Glu Trp Ser Pro Ser Thr Lys Trp His Asn Ser Tyr 200

AGG GAG CCC TTC GAG CAG CAC CTC CTG CTG GGC CTC AGC GGT TCC 645  
Arg Glu Pro Phe Glu Gln His Leu Leu Leu Gly Val Ser Val Ser 215

TGC ATT GTC ATC CTC CCC GTC TGC CTG TTG TGC TAT GTC AGC ATC 690  
Cys Ile Val Ile Leu Ala Val Cys Leu Leu Cys Tyr Val Ser Ile 230

ACC AAG ATT AAG AAA GAA TGG TGG GAT CAG ATT CCC AAC CCA GGC 735  
Thr Lys Ile Lys Lys Glu Trp Trp Asp Gln Ile Pro Asn Pro Ala 245

(57) Abstract: Methods for treating medical conditions induced by interleukin-4 involve administering an IL-4 antagonist to a patient afflicted with such a condition. Suitable IL-4 antagonists include, but are not limited to, IL-4 receptors (such as a soluble human IL-4 receptor), antibodies that bind IL-4, antibodies that bind IL-4R, IL-4 muteins that bind to IL-4R but do not induce a biological response, molecules that inhibit IL-4-induced signal transduction, and other compounds that inhibit a biological effect that results from the binding of IL-4 to a cell surface IL-4R. Particular antibodies provided herein include human monoclonal antibodies generated by procedures involving immunization of transgenic mice. Such human antibodies may be raised against human IL-4 receptor. Certain of the antibodies inhibit both IL-4-induced and IL-13-induced biological activities.

WO 01/92340 A3



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:

25 April 2002

**Published:**

— with international search report

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/17094

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K16/28 A61K39/395 A61P19/02 A61P31/04 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EMBASE, WPI Data, PAJ, EP0-Internal, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 33735 A (CELL GENESYS INC.) 31 October 1996 (1996-10-31) page 15, line 9 - line 11 claims 1,19,20,26,27 ---	1,2,4-8
X	WO 96 34096 A (CELL GENESYS INC.) 31 October 1996 (1996-10-31) page 14, line 16 - line 18 claims 1,17,18,24,25 ---	1,2,4-8
A	EP 0 604 693 A (SCHERING-PLOUGH) 6 July 1994 (1994-07-06) claims ---	1-13
A	WO 94 21282 A (IMMUNEX CORPORATION) 29 September 1994 (1994-09-29) examples claims ---	11
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

7 February 2002

Date of mailing of the international search report

26/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Nooi, F



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/17094

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>O. HULTGREN ET AL.: "Staphylococcus aureus-induced septic arthritis and septic death is decreased in IL-4-deficient mice: role of IL-4 as promoter for bacterial growth." THE JOURNAL OF IMMUNOLOGY, vol. 160, no. 10, 15 May 1998 (1998-05-15), pages 5082-5087, XP002189479 Baltimore, MD, USA abstract ---</p>	9-13
A	<p>S. ZURAWSKI ET AL.: "The primary binding subunit of the human interleukin-4 receptor is also a component of the interleukin-13 receptor." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 23, 9 June 1995 (1995-06-09), pages 13869-13878, XP000986235 Baltimore, MD, USA abstract ---</p>	1,2
A	<p>WO 91 09059 A (SCHERING CORPORATION) 27 June 1991 (1991-06-27) page 17, line 12 -page 18, line 3 examples claims ---</p>	1,12
A	<p>WO 00 18932 A (REGENERON PHARMACEUTICALS INC.) 6 April 2000 (2000-04-06) examples 10-13 figures 21-23 -----</p>	11

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 9,12,13 (all partially)

Present claims 9, 12 and 13 relate to a compound defined by reference to a desirable characteristic or property, namely an IL-4 antagonist. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to blocking monoclonal antibodies against IL-4 as disclosed in examples 1, 4-6, 8 and 9.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

**FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-8, 10, 12 (completely), 9, 13 (both partially)

Antibodies against the IL-4R, including human monoclonal antibodies against the human IL-4R, which can inhibit an IL-4-induced biological activity, including inhibition of binding of IL-4 to the IL-4R. Method for in vivo use and treatment, including that of septic arthritis.

2. Claims: 9 and 13 (both partially), 11 (completely)

Method of treatment of septic arthritis in a mammal by an IL-4 antagonist, wherein said antagonist is the soluble form of the human IL-4R of seq. id. no. 2, and the mammal is a human.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 01/17094

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9633735	A	31-10-1996	AU 5632296 A	18-11-1996
			CA 2219361 A1	31-10-1996
			EP 0822830 A1	11-02-1998
			JP 11505523 T	21-05-1999
			WO 9633735 A1	31-10-1996
			US 6075181 A	13-06-2000
			US 6150584 A	21-11-2000
WO 9634096	A	31-10-1996	CA 2219486 A1	31-10-1996
			WO 9634096 A1	31-10-1996
			AU 2466895 A	18-11-1996
			EP 0823941 A1	18-02-1998
			JP 11505107 T	18-05-1999
EP 604693	A	06-07-1994	WO 9414975 A1	07-07-1994
			EP 0604693 A1	06-07-1994
			AU 5852094 A	19-07-1994
WO 9421282	A	29-09-1994	AU 696560 B2	10-09-1998
			AU 6367794 A	11-10-1994
			EP 0689451 A1	03-01-1996
			JP 8508035 T	27-08-1996
			NZ 263102 A	26-05-1997
			WO 9421282 A1	29-09-1994
			US 6063371 A	16-05-2000
WO 9109059	A	27-06-1991	AT 136906 T	15-05-1996
			AU 639754 B2	05-08-1993
			AU 7176291 A	18-07-1991
			CA 2071908 A1	21-06-1991
			CZ 9006352 A3	13-08-1997
			DE 69026627 D1	23-05-1996
			DE 69026627 T2	22-08-1996
			DK 506826 T3	13-05-1996
			EP 0506826 A1	07-10-1992
			ES 2085983 T3	16-06-1996
			FI 922847 A	18-06-1992
			GR 3020325 T3	30-09-1996
			HU 215245 B	30-11-1998
			HU 64568 A2	28-01-1994
			IE 904589 A1	03-07-1991
			JP 7030116 B	05-04-1995
			JP 4506359 T	05-11-1992
			KR 9616862 B1	23-12-1996
			MX 9203401 A1	01-07-1992
			NO 922457 A	19-06-1992
			NZ 236511 A	25-02-1993
			OA 9704 A	30-08-1993
			PT 96230 A , B	15-10-1991
			SK 635290 A3	11-06-1999
			WO 9109059 A1	27-06-1991
			ZA 9010190 A	28-08-1991
WO 0018932	A	06-04-2000	AU 6499499 A	17-04-2000
			EP 1115876 A2	18-07-2001
			NO 20011513 A	25-05-2001
			WO 0018932 A2	06-04-2000